 An official website of the United States government
[Here's how you know](#)



MENU

Search EPA.gov


Climate Change Indicators

CONTACT US <<https://epa.gov/climate-indicators/forms/contact-us-about-climate-indicators>>

Climate Change Indicators: Atmospheric Concentrations of Greenhouse Gases

This indicator describes how the levels of major greenhouse gases in the atmosphere have changed over time.

Figure 1. Global Atmospheric Concentrations of Carbon Dioxide Over Time



<https://epa.gov/sites/default/files/2021-03/ghg-concentrations_fig-1.csv>



<https://epa.gov/sites/default/files/2021-03/ghg-concentrations_download1_2021.png>



<<https://edap.epa.gov/public/extensions/ccdataviewer/ccdataviewer.html#greenhouse-gases&atm-conc-ghg>>

This figure shows concentrations of carbon dioxide in the atmosphere from hundreds of thousands of years ago through 2019, measured in parts per million (ppm). The data come from a variety of historical ice core studies and recent air monitoring sites around the world. Each line represents a different data source.




Figure 1. Global Atmospheric Concentrations of Carbon Dioxide Over Time




Figure 2. Global Atmospheric Concentrations of Methane Over Time




Figure 3. Global Atmospheric Concentrations of Nitrous Oxide Over Time




Figure 4. Global Atmospheric Concentrations of Selected Halogenated Gases, 1978–2018






Figure 5. Global Atmospheric Concentrations of Ozone, 1979–2018

Key Points | Background | About the Indicator | About the Data | Technical Documentation

Key Points

- Global atmospheric concentrations of carbon dioxide, methane, nitrous oxide, and certain manufactured greenhouse gases have all risen significantly over the last few hundred years (see Figures 1, 2, 3, and 4).
- Historical measurements show that the current global atmospheric concentrations of carbon dioxide, methane, and nitrous oxide are unprecedented compared with the past 800,000 years (see Figures 1, 2, and 3).
- Carbon dioxide concentrations have increased substantially since the beginning of the industrial era, rising from an annual average of 280 ppm in the late 1700s to 410 ppm in 2019 (average of five sites in Figure 1)—a 46 percent increase. Almost all of this increase is due to human activities.¹
- The concentration of methane in the atmosphere has more than doubled since preindustrial times, reaching over 1,800 ppb in recent years (see the range of measurements for 2019 in Figure 2). This increase is predominantly due to agriculture and fossil fuel use.²
- Over the past 800,000 years, concentrations of nitrous oxide in the atmosphere rarely exceeded 280 ppb. Levels have risen since the 1920s, however, reaching a new high of 331 ppb in 2019 (average of three sites in Figure 3). This increase is primarily due to agriculture.³

- Concentrations of many of the halogenated gases shown in Figure 4 were essentially zero a few decades ago but have increased rapidly as they have been incorporated into industrial products and processes. Some of these chemicals have been or are currently being phased out of use because they are ozone-depleting substances, meaning they also cause harm to the Earth’s protective ozone layer. As a result, concentrations of many major ozone-depleting gases have begun to stabilize or decline (see Figure 4, left panel). Concentrations of other halogenated gases have continued to rise, however, especially where the gases have emerged as substitutes for ozone-depleting chemicals (see Figure 4, right panel).
- Overall, the total amount of ozone in the atmosphere decreased by about 3 percent between 1979 and 2018 (see Figure 5). All of the decrease happened in the stratosphere, with most of the decrease occurring between 1979 and 1994. Changes in stratospheric ozone reflect the effect of ozone-depleting substances. These chemicals have been released into the air for many years, but recently, international efforts have reduced emissions and phased out their use.
- Globally, the amount of ozone in the troposphere increased by about 13 percent between 1979 and 2018 (see Figure 5).

Water Vapor as a Greenhouse Gas

Water vapor is the most abundant greenhouse gas in the atmosphere. Human activities have only a small direct influence on atmospheric concentrations of water vapor, primarily through irrigation and deforestation, so it is not included in this indicator.⁴ The surface warming caused by human production of other greenhouse gases, however, leads to an increase in atmospheric water vapor because warmer temperatures make it easier for water to evaporate and stay in the air in vapor form. This creates a positive “feedback loop” in which warming leads to more warming.

Background

Since the Industrial Revolution began in the 1700s, people have added a substantial amount of greenhouse gases into the atmosphere by burning fossil fuels, cutting down forests, and conducting other activities (see the U.S. <https://epa.gov/climate-indicators/climate-change-indicators-us-greenhouse-gas-emissions> and Global <https://epa.gov/climate-indicators/climate-change-indicators-global-greenhouse-gas-emissions> Greenhouse Gas Emissions indicators). When greenhouse gases are emitted into the atmosphere, many remain there for long time periods ranging from a decade to many millennia. Over time, these gases are removed from the atmosphere by chemical reactions or by emissions sinks, such as the oceans and vegetation, which absorb greenhouse gases from the atmosphere. As a result of human activities, however, these gases are entering the atmosphere more quickly than they are being removed, and thus their concentrations are increasing.

Carbon dioxide, methane, nitrous oxide, and certain manufactured gases called halogenated gases (gases that contain chlorine, fluorine, or bromine) become well mixed throughout the global atmosphere because of their relatively long lifetimes and because of transport by winds. Concentrations of these greenhouse gases are measured in parts per million (ppm), parts per billion (ppb), or parts per trillion (ppt) by volume. In other words, a concentration of 1 ppb for a given gas means there is one molecule of that gas in every 1 billion molecules of air. Some halogenated gases are considered major greenhouse gases due to their very high global warming potentials and long atmospheric lifetimes even if they only exist at a few ppt (see table <https://epa.gov/climate-indicators/greenhouse-gases#major-long-lived-greenhouse-gases-and-their-characteristics>).

This indicator looks at global average levels of ozone in both the stratosphere and troposphere. For trends in ground-level ozone concentrations within the United States, see EPA's *National Air Quality Trends Report* at: www.epa.gov/air-trends <https://epa.gov/air-trends>.

Ozone is also a greenhouse gas, but it differs from other greenhouse gases in several ways. The effects of ozone depend on its altitude, or where the gas is located vertically in the atmosphere. Most ozone naturally exists in the layer of the atmosphere called the stratosphere, which ranges from approximately 6 to 30 miles above the Earth's surface. Ozone in the stratosphere has a slight net warming effect on the planet, but it is good

for life on Earth because it absorbs harmful ultraviolet radiation from the sun, preventing it from reaching the Earth's surface. In the troposphere—the layer of the atmosphere near ground level—ozone is an air pollutant that is harmful to breathe, a main ingredient of urban smog, and an important greenhouse gas that contributes to climate change (see the Climate Forcing <<https://epa.gov/climate-indicators/climate-change-indicators-climate-forcing>> indicator). Unlike the other major greenhouse gases, tropospheric ozone only lasts for days to weeks, so levels often vary by location and by season.

About the Indicator

This indicator describes concentrations of greenhouse gases in the atmosphere. It focuses on the major greenhouse gases that result from human activities.

For carbon dioxide, methane, nitrous oxide, and halogenated gases, recent measurements come from monitoring stations around the world, while measurements of older air come from air bubbles trapped in layers of ice from Antarctica and Greenland. By determining the age of the ice layers and the concentrations of gases trapped inside, scientists can learn what the atmosphere was like thousands of years ago.

This indicator also shows data from satellite instruments that measure ozone density in the troposphere, the stratosphere, and the “total column,” or all layers of the atmosphere. These satellite data are routinely compared with ground-based instruments to confirm their accuracy. Ozone data have been averaged worldwide for each year to smooth out the regional and seasonal variations.

About the Data

Indicator Notes

This indicator includes several of the most important halogenated gases, but some others are not shown. Many other halogenated gases are also greenhouse gases, but Figure 4 is limited to a set of common examples that represent most of the major types of these gases. The indicator also does not address certain other pollutants that can affect climate by either reflecting or absorbing energy. For example, sulfate particles can reflect sunlight away from the Earth, while black carbon aerosols (soot) absorb energy. Data for nitrogen trifluoride (Figure 4) reflect modeled averages based on

measurements made in the Northern Hemisphere and some locations in the Southern Hemisphere, to represent global average concentrations over time. The global averages for ozone only cover the area between 50°N and 50°S latitude (77 percent of the Earth's surface), because at higher latitudes the lack of sunlight in winter creates data gaps and the angle of incoming sunlight during the rest of the year reduces the accuracy of the satellite measuring technique.

Data Sources

Global atmospheric concentration measurements for carbon dioxide (Figure 1), methane (Figure 2), and nitrous oxide (Figure 3) come from a variety of monitoring programs and studies published in peer-reviewed literature. Global atmospheric concentration data for selected halogenated gases (Figure 4) were compiled by the Advanced Global Atmospheric Gases Experiment and the National Oceanic and Atmospheric Administration. A similar figure with many of these gases appears in the Intergovernmental Panel on Climate Change's Fifth Assessment Report.¹⁴ Satellite measurements of ozone were processed by the National Aeronautics and Space Administration and validated using ground-based measurements collected by the National Oceanic and Atmospheric Administration.

Technical Documentation

- Download related technical information PDF <<https://epa.gov/climate-indicators/downloads-indicators-technical-documentation>>

References

¹ USGCRP (U.S. Global Change Research Program). 2017. Climate science special report: Fourth National Climate Assessment, volume I. Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock, eds. <https://science2017.globalchange.gov> <<https://science2017.globalchange.gov>>. doi:10.7930/J0J964J6.

² IPCC (Intergovernmental Panel on Climate Change). 2013. Climate change 2013: The physical science basis. Working Group I contribution to the IPCC Fifth Assessment Report. Cambridge, United Kingdom: Cambridge University Press. www.ipcc.ch/report/ar5/wg1 EXIT <<https://www.ipcc.ch/report/ar5/wg1>>.

3. USGCRP (U.S. Global Change Research Program). 2017. Climate science special report: Fourth National Climate Assessment, volume I. Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock, eds. <https://science2017.globalchange.gov> <<https://science2017.globalchange.gov>>. doi:10.7930/J0J964J6.
4. USGCRP (U.S. Global Change Research Program). 2017. Climate science special report: Fourth National Climate Assessment, volume I. Wuebbles, D.J., D.W. Fahey, K.A. Hibbard, D.J. Dokken, B.C. Stewart, and T.K. Maycock, eds. <https://science2017.globalchange.gov> <<https://science2017.globalchange.gov>>. doi:10.7930/J0J964J6.
5. [see full list provided below]
6. [see full list provided below]
7. [see full list provided below]
8. AGAGE (Advanced Global Atmospheric Gases Experiment). 2019. ALE/GAGE/AGAGE database. Updated January 8, 2019. Accessed December 2020. http://agage.eas.gatech.edu/data_archive/global_mean EXIT <http://agage2.eas.gatech.edu/data_archive/global_mean>.
9. NOAA (National Oceanic and Atmospheric Administration). 2019. Halocarbons and Other Atmospheric Trace Species group (HATS). Updated October 2019. Accessed January 2021. ftp://ftp.cmdl.noaa.gov/hats/Total_Cl_Br <ftp://ftp.cmdl.noaa.gov/hats/total_cl_br%e2%80%8b>.
10. Rigby, M. 2017 update to data originally published in: Arnold, T., C.M. Harth, J. Mühle, A.J. Manning, P.K. Salameh, J. Kim, D.J. Ivy, L.P. Steele, V.V. Petrenko, J.P. Severinghaus, D. Baggenstos, and R.F. Weiss. 2013. Nitrogen trifluoride global emissions estimated from updated atmospheric measurements. P. Natl. Acad. Sci. USA 110(6):2029–2034. Data updated December 2017.
11. NASA (National Aeronautics and Space Administration). 2013. Data—TOMS/SBUV TOR data products. Accessed November 2013. <https://science-data.larc.nasa.gov/TOR/data.html> <<https://science-data.larc.nasa.gov/tor/data.html>>.
12. NASA (National Aeronautics and Space Administration). 2019. SBUV merged ozone data set (MOD). Version 8.6. Updated November 20, 2019. Accessed December 2020. https://acd-ext.gsfc.nasa.gov/Data_services/merged/index.html <https://acd-ext.gsfc.nasa.gov/data_services/merged/index.html>

¹³. NASA (National Aeronautics and Space Administration). 2020. Tropospheric ozone data from AURA OMI/MLS. Accessed December 2020. http://acdb-ext.gsfc.nasa.gov/Data_services/cloud_slice/new_data.html <http://acdb-ext.gsfc.nasa.gov/data_services/cloud_slice/new_data.html>.

¹⁴. IPCC (Intergovernmental Panel on Climate Change). 2013. Climate change 2013: The physical science basis. Working Group I contribution to the IPCC Fifth Assessment Report. Cambridge, United Kingdom: Cambridge University Press. www.ipcc.ch/report/ar5/wg1 EXIT <<https://www.ipcc.ch/report/ar5/wg1%e2%80%8b>>.

Atmospheric Concentrations of Greenhouse Gases: Citations for Figures 1, 2, and 3

FIGURE 1

Antarctic Ice Cores: approximately 803,719 BCE to 2001 CE

Bereiter, B., S. Eggleston, J. Schmitt, C. Nehrbass-Ahles, T.F. Stocker, H. Fischer, S. Kipfstuhl, and J. Chappellaz. 2015. Revision of the EPICA Dome C CO₂ record from 800 to 600 kyr before present. *Geophys. Res. Lett.* 42(2):542–549. www.ncdc.noaa.gov/paleo-search/study/17975 <<https://www.ncdc.noaa.gov/paleo-search/study/17975>>.

Mauna Loa, Hawaii: 1959 CE to 2019 CE

NOAA (National Oceanic and Atmospheric Administration). 2020. Annual mean carbon dioxide concentrations for Mauna Loa, Hawaii. Updated September 23, 2020. Accessed December 29, 2020. ftp://ftp.cmdl.noaa.gov/products/trends/co2/co2_annmean_mlo.txt <ftp://ftp.cmdl.noaa.gov/products/trends/co2/co2_annmean_mlo.txt>.

Barrow, Alaska: 1974 CE to 2019 CE

Cape Matatula, American Samoa: 1976 CE to 2019 CE

South Pole, Antarctica: 1976 CE to 2019 CE

NOAA (National Oceanic and Atmospheric Administration). 2020. Monthly mean carbon dioxide concentrations for Barrow, Alaska; Cape Matatula, American Samoa; and the South Pole. Updated August 26, 2020. Accessed December 29, 2020. ftp://ftp.cmdl.noaa.gov/data/trace_gases/co2/in-situ/surface <ftp://ftp.cmdl.noaa.gov/data/trace_gases/co2/in-situ/surface>.

Cape Grim, Australia: 1977 CE to 2019 CE

CSIRO (Commonwealth Scientific and Industrial Research Organisation). 2020. Monthly mean baseline (background) carbon dioxide concentrations measured at the Cape Grim Baseline Air Pollution Station, Tasmania, Australia. Updated December 2020. Accessed December 29, 2020.

http://capegrim.csiro.au/GreenhouseGas/data/CapeGrim_CO2_data_download.csv EXIT <http://capegrim.csiro.au/greenhousegas/data/capegrim_co2_data_download.csv>.

Shetland Islands, Scotland: 1993 CE to 2002 CE

Steele, L.P., P.B. Krummel, and R.L. Langenfelds. 2007. Atmospheric CO₂ concentrations (ppmv) derived from flask air samples collected at Cape Grim, Australia, and Shetland Islands, Scotland. Commonwealth Scientific and Industrial Research Organisation. Accessed January 20, 2009.

<http://cdiac.esd.ornl.gov/ftp/trends/co2/csiro> <<http://cdiac.esd.ornl.gov/ftp/trends/co2/csiro>>.

Lampedusa Island, Italy: 1993 CE to 2000 CE

Chamard, P., L. Ciattaglia, A. di Sarra, and F. Monteleone. 2001. Atmospheric carbon dioxide record from flask measurements at Lampedusa Island. In: Trends: A compendium of data on global change. Oak Ridge, TN: U.S. Department of Energy. Accessed September 14, 2005. <http://cdiac.ess-dive.lbl.gov/trends/co2/lampis.html>

<<http://cdiac.ess-dive.lbl.gov/trends/co2/lampis.html>>

FIGURE 2

EPICA Dome C, Antarctica: approximately 797,446 BCE to 1937 CE

Loulergue, L., A. Schilt, R. Spahni, V. Masson-Delmotte, T. Blunier, B. Lemieux, J.-M. Barnola, D. Raynaud, T.F. Stocker, and J. Chappellaz. 2008. Orbital and millennial-scale features of atmospheric CH₄ over the past 800,000 years. *Nature* 453:383–386. www.ncdc.noaa.gov/paleo-search/study/6093

<<https://www.ncdc.noaa.gov/paleo-search/study/6093>>.

Law Dome, Antarctica: approximately 1008 CE to 1980 CE

Etheridge, D.M., L.P. Steele, R.J. Francey, and R.L. Langenfelds. 2002. Historic CH₄ records from Antarctic and Greenland ice cores, Antarctic firn data, and archived air samples from Cape Grim, Tasmania. In: Trends: A compendium of data on global change. Oak Ridge, TN: U.S. Department of Energy. Accessed September 13, 2005. http://cdiac.ess-dive.lbl.gov/trends/atm_meth/lawdome_meth.html <http://cdiac.ess-dive.lbl.gov/trends/atm_meth/lawdome_meth.html>.

Cape Grim, Australia: 1985 CE to 2019 CE

CSIRO (Commonwealth Scientific and Industrial Research Organisation). 2020. Monthly mean baseline (background) methane concentrations measured at the Cape Grim Baseline Air Pollution Station, Tasmania, Australia. Updated December 2020. Accessed December 29, 2020.

http://capegrim.csiro.au/GreenhouseGas/data/CapeGrim_CH4_data_download.csv EXIT

<http://capegrim.csiro.au/greenhousegas/data/capegrim_ch4_data_download.csv>.

Mauna Loa, Hawaii: 1984 CE to 2019 CE

NOAA (National Oceanic and Atmospheric Administration). 2020. Monthly mean CH₄ concentrations for Mauna Loa, Hawaii. Updated July 24, 2020. Accessed December 29, 2020.

ftp://ftp.cmdl.noaa.gov/data/trace_gases/ch4/flask/surface/ch4_mlo_surface-flask_1_ccgg_month.txt
<ftp://ftp.cmdl.noaa.gov/data/trace_gases/ch4/flask/surface/ch4_mlo_surface-flask_1_ccgg_month.txt>.

Shetland Islands, Scotland: 1993 CE to 2001 CE

Steele, L.P., P.B. Krummel, and R.L. Langenfelds. 2002. Atmospheric methane record from Shetland Islands, Scotland (October 2002 version). In: Trends: A compendium of data on global change. Oak Ridge, TN: U.S. Department of Energy. Accessed September 13, 2005. http://cdiac.ess-dive.lbl.gov/trends/atm_meth/csiro/csiro-shetlandch4.html <http://cdiac.ess-dive.lbl.gov/trends/atm_meth/csiro/csiro-shetlandch4.html>.

FIGURE 3

EPICA Dome C, Antarctica: approximately 796,475 BCE to 1937 CE

Schilt, A., M. Baumgartner, T. Blunier, J. Schwander, R. Spahni, H. Fischer, and T.F. Stocker. 2010. Glacial-interglacial and millennial scale variations in the atmospheric nitrous oxide concentration during the last 800,000 years. *Quaternary Sci. Rev.* 29:182–192. www.ncdc.noaa.gov/paleo-search/study/8615
<<http://www.ncdc.noaa.gov/paleo-search/study/8615>>.

Antarctica: approximately 1903 CE to 1976 CE

Battle, M., M. Bender, T. Sowers, P. Tans, J. Butler, J. Elkins, J. Ellis, T. Conway, N. Zhang, P. Lang, and A. Clarke. 1996. Atmospheric gas concentrations over the past century measured in air from firn at the South Pole. *Nature* 383:231–235. Data available at: https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=797.

Cape Grim, Australia: 1979 CE to 2019 CE

CSIRO (Commonwealth Scientific and Industrial Research Organisation). 2020c. Monthly mean baseline (background) nitrous oxide concentrations measured at the Cape Grim Baseline Air Pollution Station, Tasmania, Australia. Updated December 2020. Accessed December 29, 2020. http://capegrim.csiro.au/GreenhouseGas/data/CapeGrim_N2O_data_download.csv EXIT
<http://capegrim.csiro.au/greenhousegas/data/capegrim_n2o_data_download.csv>.

South Pole, Antarctica: 1998 CE to 2019 CE

Barrow, Alaska: 1999 CE to 2019 CE

Mauna Loa, Hawaii: 2000 CE to 2019 CE

NOAA (National Oceanic and Atmospheric Administration). 2020. Monthly mean N2O concentrations for Barrow, Alaska; Mauna Loa, Hawaii; and the South Pole. Accessed December 29, 2020. www.esrl.noaa.gov/gmd/hats/insitu/cats/cats_conc.html
<https://www.esrl.noaa.gov/gmd/hats/insitu/cats/cats_conc.html>.

Learn about other indicators in this section



<[https://epa.gov/climate-indicators/climate-change-indicators-us-greenhouse-](https://epa.gov/climate-indicators/climate-change-indicators-us-greenhouse-gas-emissions)

gas-emissions>



<[https://epa.gov/climate-indicators/climate-change-indicators-](https://epa.gov/climate-indicators/climate-change-indicators-global-greenhouse-gas-emissions)

global-greenhouse-gas-emissions>

<[https://epa.gov/climate-indicators/climate-](https://epa.gov/climate-indicators/climate-change-indicators-atmospheric-concentrations-greenhouse-gases)

change-indicators-atmospheric-concentrations-greenhouse-gases>



<<https://epa.gov/climate-indicators/climate-change-indicators-climate-forcing>>

[Climate Change Indicators <https://epa.gov/climate-indicators>](https://epa.gov/climate-indicators)

Greenhouse Gases

U.S. Greenhouse Gas Emissions <<https://epa.gov/climate-indicators/climate-change-indicators-us-greenhouse-gas-emissions>>

Global Greenhouse Gas Emissions <<https://epa.gov/climate-indicators/climate-change-indicators-global-greenhouse-gas-emissions>>

Atmospheric Concentrations of Greenhouse Gases

Climate Forcing <<https://epa.gov/climate-indicators/climate-change-indicators-climate-forcing>>

Weather and Climate <<https://epa.gov/climate-indicators/weather-climate>>

Oceans <<https://epa.gov/climate-indicators/oceans>>

Snow and Ice <<https://epa.gov/climate-indicators/snow-ice>>

Health and Society <<https://epa.gov/climate-indicators/health-society>>

Ecosystems <<https://epa.gov/climate-indicators/ecosystems>>

Frequent Questions <<https://epa.gov/climate-indicators/frequent-questions-about-climate-change-indicators>>

Contact Us <<https://epa.gov/climate-indicators/forms/contact-us-about-climate-indicators>> to ask a question, provide feedback, or report a problem.



Discover.

Accessibility <<https://epa.gov/accessibility>>

Related Documents

Budget & Performance <<https://epa.gov/planandbudget>>

Contracting <<https://epa.gov/contracts>>

EPA www Web Snapshot <<https://epa.gov/home/wwwepagov-snapshots>>

Grants <<https://epa.gov/grants>>

No FEAR Act Data <<https://epa.gov/ocr/whistleblower-protections-epa-and-how-they-relate-non-disclosure-agreements-signed-epa-employees>>

Plain Writing <<https://epa.gov/web-policies-and-procedures/plain-writing>>

Privacy <<https://epa.gov/privacy>>

Privacy and Security Notice <<https://epa.gov/privacy/privacy-and-security-notice>>

Connect.

Data.gov <<https://www.data.gov/>>

Inspector General <<https://epa.gov/office-inspector-general/about-epas-office-inspector-general>>

Jobs <<https://epa.gov/careers>>

Newsroom <<https://epa.gov/newsroom>>

Open Government <<https://epa.gov/data>>

Regulations.gov <<https://www.regulations.gov/>>

Subscribe <<https://epa.gov/newsroom/email-subscriptions-epa-news-releases>>

USA.gov <<https://www.usa.gov/>>

White House <<https://www.whitehouse.gov/>>

Ask.

Contact EPA <<https://epa.gov/home/forms/contact-epa>>

EPA Disclaimers <<https://epa.gov/web-policies-and-procedures/epa-disclaimers>>

Hotlines <<https://epa.gov/aboutepa/epa-hotlines>>

FOIA Requests <<https://epa.gov/foia>>

Frequent Questions <<https://epa.gov/home/frequent-questions-specific-epa-programtopics>>

Follow.



LAST UPDATED ON JULY 21, 2021