



Rincon Consultants, Inc.

2215 Faraday Avenue  
Suite A  
Carlsbad, California 92008  
760-918-9444

June 18, 2023  
Project No: 22-13460

Stacey Love  
U.S. Fish and Wildlife Service, Carlsbad Office  
2177 Salk Avenue #250  
Carlsbad, California 92008  
Via email: [stacey\\_love@fws.gov](mailto:stacey_love@fws.gov)

**Subject: Coastal California Gnatcatcher Protocol Survey Report  
Woodward Specific Plan Development Project  
San Marcos, San Diego County, California**

Dear Ms. Love:

Rincon conducted protocol breeding season coastal California gnatcatcher (*Poliioptila californica californica*; CAGN) for the proposed Woodward Specific Plan Development Project (Project) to further evaluate presence/absence of the species as part of the environmental review process. Rincon is submitting this letter report summarizing the results of focused surveys for the federally threatened CAGN.

Surveys were conducted by Rincon biologist Kelly Rios who currently holds an Endangered and Threatened Species Permit issued by the United States (U.S.) Fish and Wildlife Service (USFWS), Permit TE 018909-6, under Section 10(a)(1)(A) of the Federal Endangered Species Act. The 15-day notification letter of intent to conduct protocol breeding season surveys for CAGN was sent to the USFWS Carlsbad office on April 18, 2023.


## **Project Location and Environmental Setting**

The 8.57-acre project site is located in the city of San Marcos and is situated 0.4 mile north of Highway 78 (Figure 1). The project site is located within the U.S. Geological Survey 7.5-minute topographic quadrangle San Marcos, California (See attached map, Figure 2). The project site is specifically located to the east of Woodward Street and north of East Mission Road (Figure 3). The project site is also located within the boundaries of the San Diego County Multiple Habitat Conservation Program (MHCP) subarea plan and is located adjacent to a 7.73-acre open space MHCP Hardline Reserve land to the north, east, and west of the project site across Woodward Street (Figure 4).

The City of San Marcos implements portions of the MHCP through subarea plans, which describe specific implementing mechanisms such as the MHPA pursuant to Section 10(a) of the Federal ESA and the Natural Community Conservation Program (NCCP) plan pursuant to the California NCCP Act of 1991 and the California ESA.

Focused CAGN surveys were conducted for 16.46 acres of Diegan coastal sage scrub and disturbed Diegan coastal sage scrub habitat within the project site and surrounding 250 foot buffer, referred to as the "study area/survey area" (Figure 5). The project site occurs on a rolling hillside with slopes generally facing west and southwest.


**Figure 1 Regional Location**



Imagery provided by Esri and its licensors © 2023.


22-133460 Bio Figures  
Fig 1 Regional Location

Project Location





**Figure 2 Project Location on USGS 7.5-minute Topographic Quadrangle**



Basemap provided by National Geographic Society, Esri, and their licensors © 2023. San Marcos Quadrangle, T12S R03W S11.  
The topographic representation depicted in this map may not portray all of the features currently found in the vicinity today and/or features depicted in this map may have changed since the original topographic map was assembled.



Figure 3 Project Site





**Figure 4 Project Vicinity to MHCP Conservation Areas**



Imagery provided by Microsoft Bing and its licensors © 2023.  
 Additional data provided by SANDAG, 2018.

22-133460 Bio Figures  
 Fig X Project Vicinity to MHCP Conservation Areas



**Figure 5 Project Site, Project Impact Area, and Vegetation Map**



Imagery provided by Microsoft Bing and its licensors © 2023.

22-133460 Bio Figures  
 Fig 3 Vegetation 20230426



## Diegan Coastal Sage Scrub

Diegan Coastal Sage Scrub is dominated by low, soft-woody subshrubs that are most active in winter and early spring. Many taxa are facultatively drought-deciduous and typically grow in low moisture-availability sites, such as steep, xeric slopes or clay-rich soils that are slow to release stored water. Dominant plant species in Diegan Coastal Sage Scrub usually include California sagebrush (*Artemisia californica*), California buckwheat (*Eriogonum fasciculatum*), deerweed (*Acmispon glaber*), chaparral mallow (*Malacothamnus fasciculatus*), laurel sumac (*Malosma laurina*), lemonade berry (*Rhus integrifolia*), and black sage (*Salvia mellifera*) (Oberbauer et al. 2008). Diegan Coastal Sage Scrub is considered a sensitive community by the City, falling under the Habitat Group C.

Diegan Coastal Sage Scrub is dominated by low, soft-woody subshrubs that are most active in winter and early spring. Many taxa are facultatively drought-deciduous. This community is typically on low moisture-availability sites such as steep, xeric slopes or clay-rich soils that are slow to release stored water.

This vegetation community is the most prevalent community within the study area (16.27 acres). The majority of the project site, as well as the study area to the north, supports high-quality Diegan Coastal Sage Scrub.

The shrub layer is dense and dominated by laurel sumac (*Malosma laurina*), California sagebrush (*Artemisia californica*) and lemonade berry (*Rhus integrifolia*), with California buckwheat (*Eriogonum fasciculatum*), deerweed (*Acmispon glaber*), white sage (*Salvia apiana*), black sage (*S. mellifera*) and coyote brush (*Baccharis pilularis*) (Oberbauer et al. 2008). Several non-native species including tocalote (*Centaurea melitensis*), mustards (*brassica* sp.), slender wild oat (*Avena barbata*) and ripgut brome (*Bromus dianrus*) are also common throughout this community. The northern portion of the survey area consisted of more dense shrub cover compared to the central and southern portions.

Historical disturbance within the central portion of the site identified 0.88 acre of a more Disturbed Diegan Coastal Sage Scrub community, with evidence of revegetation in the last several years. Disturbed Diegan Coastal Sage Scrub is also present within the central and southern edges of the project site and within the study area to the south.

The central and southern portion of the project site contained disturbed Diegan Coastal Sage Scrub that was revegetating in historically disturbed areas, including fuel medication breaks. Dominant shrub species included California buckwheat, California sagebrush, black sage, coyote brush, with golden yarrow (*Eriophyllum confertiflorum*) and sparse herbaceous species, including giant woollystar (*Eriastrum densifolium*) and small seed sandmat (*Euphorbia polycarpa*) and large patches of open/bare ground.

## Species Background

The CAGN is a non-migratory songbird found on the coastal slopes of southern California. It ranges from Ventura County south to northwest Baja California, Mexico (Atwood et al. 1999; Jones and Ramirez 1995). The CAGN belongs to the old-world warbler and gnatcatcher family, *Sylviidae*. It is a small blue-gray songbird which measures 4.5 inches (11 centimeters) and weighs 0.2 ounces (6 grams). It has dark blue-gray feathers on its back and grayish-white feathers on its underside. The wings have a brownish wash. Its long tail is mostly black with white outer tail feathers. The species has a thin, small bill. The males have a black cap during the spring and summer which is absent during the winter. Both males and females have a white ring around their eyes.



The breeding season of the CAGN extends from late February through August with the peak of nesting occurring from mid-March through mid-May. The breeding territory size of the CAGN ranges from two to 22 acres (one to nine hectares), with home ranges expanding up to 39 acres (16 hectares), during the non-breeding season (Bontrager 1991; USFWS 1993). Nest parasitism by brown-headed cowbirds (*Molothrus ater*) has been documented (Unitt 1984). Typically, there is a high rate of nest failure each breeding season. This is offset by rapid and persistent re-nesting efforts; a breeding pair may attempt to nest as many as ten times in a year, producing up to three successful broods in a season (Atwood and Bontrager 2001). There is evidence that this species is also susceptible to nest predation by various animals such as snakes, coyotes (*Canis latrans*), fox, rodents, and other birds, such as California scrub-jays (*Aphelocoma californica*) (Atwood et al. 1999).

CAGN are strongly associated with coastal sage scrub habitats below 820 feet in coastal areas and between 820 and 1,640 feet in inland areas (Atwood and Bolsinger 1992); however, not all types of coastal sage scrub communities are used or preferred. This species is reported to be most abundant in areas dominated by California sagebrush and California buckwheat. CAGN numbers are generally low in coastal habitats dominated by black sage, white sage (*Salvia apiana*), or lemonade berry; in inland areas, habitats dominated by black sage may be used more regularly (Atwood and Bontrager 2001).

Population estimates for the CAGN vary. Atwood (1992) estimated that 1,811 to 2,291 pairs of CAGN existed in 1992 throughout its range in southern California. In 1996, the USFWS estimated the population in San Diego County at 3,000 pairs, excluding pairs located on sites where habitat loss had already been approved (Atwood and Bontrager 2001). According to a 1999 population estimate in San Diego and other southern California counties, the USFWS estimated the population in San Diego at 1,917 pairs, Orange County at 643 pairs, Los Angeles County at 144 pairs, San Bernardino County at 27 pairs, and Ventura County at 4 pairs (Atwood and Bontrager 2001).

The CAGN is federally listed as threatened and is a California Department of Fish and Wildlife (CDFW) Species of Special Concern. The USFWS listed the CAGN as threatened (USFWS 1993) pursuant to the federal Endangered Species Act of 1973 as amended on March 30, 1993. Critical habitat was designated for CAGN on October 24, 2000 (USFWS 2007).

The primary cause of this species' decline is the cumulative loss of its coastal sage scrub vegetation community to urban and agricultural development. The USFWS has estimated that coastal sage scrub vegetation has been reduced by 70 to 90 percent of its historical extent (USFWS 1991) and little of what remains is protected in natural open space.

## **Survey Methodology**

### **Focused Surveys**

Notification to commence protocol surveys was provided via email correspondence to the USFWS on April 18, 2023. Kelly Rios, permitted Rincon biologist, conducted all CAGN surveys in accordance with the survey protocol for CAGN surveys within MHPA areas, titled *Section III of the USFWS Coastal California Gnatcatcher Presence/Absence Survey Protocol*, issued February 28, 1997, and revised July 28, 1997. The protocol requires that between February 15 and August 30, a minimum of six surveys shall be conducted at least one week apart to determine presence/absence of CAGN.

A total of six surveys were conducted between May 3 and June 14, 2023. The six surveys occurred between 0600 and 1200 hours each day. Suitable Diegan Coastal Sage Scrub occurs within the project site boundaries, north and east of the project site, and west of the project site across





Woodward Street. The study area consists of predominately Diegan Coastal Sage Scrub and disturbed Diegan Coastal Sage Scrub. Larger, more dense shrub cover within the community dominated by laurel sumacs (*Malosma laurina*) and California sage brush occurs in the northern portion of the survey area. Additional open space MHCP Reserve Habitat is adjacent to the project on the north and east sides (Figure 5).

A total of approximately 16.46 (6.6 hectares) were surveyed and did not exceed the maximum of 100 acres per survey day. Surveys were not conducted during periods of excessive or abnormal heat, wind, fog, and other inclement weather. Surveys were conducted with binoculars to aid in bird detection. The biologist slowly walked the survey area, stopping at approximate 50-foot intervals and used an audio recording of coastal CAGN vocalizations after individuals had been initially located. Recorded CAGN vocalizations were played sparingly and only if other means of detection had failed. If a CAGN was detected before playing recorded vocalizations, the recordings were not played. If CAGN is detected in response to the tape play, use of playback was discontinued immediately. If any CAGN were observed, age, sex, breeding status, and behavioral characteristics were recorded, if possible; the protocol level surveys did not include focused nest searches.

## Survey Results

Details on the date, surveyor, time, conditions, and CAGN survey results are provided in Table 1.

**Table 1 Coastal California Gnatcatcher Survey Conditions and Results**

Date	Surveyor	Survey Conditions	Survey Results
5/3/23	Kelly Rios	630-830 AM Winds 1-2 mph 100% Cloud Cover 53-54 °F	No CAGN observed
5/10/23	Kelly Rios	945-1145 AM Winds 1-2 mph 100% Cloud Cover 59-62 °F	No CAGN observed
5/17/23	Kelly Rios	1000AM-1200 PM Winds 1-2 mph 95% Cloud Cover 65-68 °F	No CAGN observed
5/24/23	Kelly Rios	1015AM-1200PM Winds 1-2 mph 100% Cloud Cover 58-60 °F	No CAGN observed
5/31/23	Kelly Rios	930-1130 AM Winds 1-2 mph 100% Cloud Cover 1-2 °F	No CAGN observed
6/14/23	Kelly Rios	0900-1100 AM Winds 1-2 mph 100% Cloud Cover 64-66 °F	No CAGN observed

During the 2023 protocol surveys, no observations of any CAGNs were recorded. CAGN surveys were conducted in 2018 and 2020 for the Mission 316 West Project (KMEA 2019) located just south of



the project site which included the Woodward Specific Plan Project 2023 survey area. CAGN observations were documented within the subject project limits. Further details regarding these observations are provided in the Conclusions section below.

Avian activity levels and diversity were generally high during the recent surveys. Common species expected to occur within coastal sage scrub and adjacent disturbed, and urban/developed habitats were observed on a regular basis. A complete list of avian species detected by sight or sound during the surveys is included in Table 2.

**Table 2 Species Observed during Protocol Level Surveys**

<b>Scientific Family, Name</b>	<b>Species</b>
<b>Accipitridae</b>	<b>Hawks, Kites, &amp; Eagles</b>
<i>Accipiter cooperii</i>	Cooper's hawk
<i>Buteo jamaicensis</i>	red-tailed hawk
<b>Aegithalidae</b>	<b>Bushtits</b>
<i>Psaltriparus minimus</i>	bushtit
<b>Apodiformes</b>	<b>Swifts</b>
<i>Aeronautes saxatalis</i>	white-throated swift
<b>Bombycillidae</b>	<b>Waxwings</b>
<i>Bombycilla cedrorum</i>	cedar waxwing
<b>Cathartidae</b>	<b>New World Vultures</b>
<i>Cathartes aura</i>	turkey vulture
<b>Columbidae</b>	<b>Pigeons &amp; Doves</b>
<i>Zenaida macroura</i>	mourning dove
<b>Corvidae</b>	<b>Ravens, Crows, Jays, &amp; Magpies</b>
<i>Aphelocoma californica</i>	California scrub-jay
<i>Corvus brachyrhynchos</i>	American crow
<i>Corvus corax</i>	common raven
<b>Emberizidae</b>	<b>Emberizids</b>
<i>Pipilo crissalis</i>	California towhee
<i>Pipilo maculatus</i>	spotted towhee
<b>Fringillidae</b>	<b>Finches</b>
<i>Haemorhous mexicanus</i>	house finch
<i>Spinus psaltria</i>	lesser goldfinch
<b>Icteridae</b>	<b>Orioles, Grackles, Cowbirds</b>
<i>Icterus cucullatus</i>	hooded oriole
<b>Mimidae</b>	<b>Mockingbirds &amp; Thrashers</b>
<i>Mimus polyglottos</i>	Northern mockingbird
<i>Toxostoma redivivum</i>	California thrasher
<b>Timaliidae</b>	<b>Babblers</b>
<i>Chamaea fasciata</i>	wrentit
<b>Troglodytidae</b>	<b>Wrens</b>
<i>Thryomanes bewickii</i>	Bewick's wren





Scientific Family, Name	Species
<i>Troglodytes aedon</i>	house wren
<b>Passerellidae</b>	<b>Sparrows</b>
<i>Melospiza melodia</i>	song sparrow
<b>Parulidae</b>	<b>New World Warblers</b>
<i>Setophaga petechia</i>	yellow warbler
<b>Picidae</b>	<b>Woodpeckers, Wrynecks, and Sapsuckers</b>
<i>Colaptes auratus</i>	Northern flicker
<i>Dryobates nuttallii</i>	Nuttall's woodpecker
<b>Odontophoridae</b>	<b>New World Quail</b>
<i>Callipepla californica</i>	California quail
<b>Trochilidae</b>	<b>Hummingbirds</b>
<i>Calypte anna</i>	Anna's hummingbird
<b>Tyrannidae</b>	<b>Tyrant Flycatchers</b>
<i>Empidonax difficilis</i>	Pacific-slope flycatcher
<i>Sayornis nigricans</i>	black phoebe
<i>Tyrannus vociferans</i>	Cassin's kingbird

## Conclusions

During the 2023 CAGN protocol surveys, no individuals, pairs, or nests were observed within the study area or adjacent areas. CAGN had been previously identified in the Diegan Coastal Sage Scrub within the project area, including within the overall study area to the east and west, during USFWS protocol surveys that were conducted in 2018 and preconstruction surveys conducted in 2020 for the adjacent Mission 316 development project located south of the project area (KMEA 2019). The Survey Area for the Mission 316 protocol CAGN surveys totaled 28 acres, which included a 500-foot buffer around the 3.7-acre project to the south. These surveys recorded individuals foraging, nesting pairs, and two family groups within the proposed Woodward project area. Individuals were also recorded within sage scrub habitat and designated MHCP Hardline Reserve to the west of the project site across Woodward Street. Foraging individuals were also observed and recorded to the adjacent area east of the project site, south of Silk Mill Place Road.

## Recommendations

Although no CAGN were observed, Rincon recommends conducting pre-disturbance protocol surveys and implementing additional avoidance measures in accordance with mitigation measures outlined in the Biological Resources Technical Report (Rincon, 2023) prepared for the City of San Marcos.

## Certification

I certify that the information in this survey report and attached exhibits fully and accurately represents my work.

Sincerely,  
**Rincon Consultants, Inc.**



Handwritten signature of Kelly Rios in black ink.

Kelly Rios, TE 018909-6  
Permitted Biologist

Handwritten signature of Jared Reed in black ink.

Jared Reed  
Project Manager/Senior Biologist

Handwritten signature of Jacob Hargis in black ink.

Jacob Hargis  
Biologist





## References

- Atwood, J. L. 1992. A maximum estimate of the California Gnatcatcher's population size in the United States. *Western Birds* 23:1-9.
- Atwood, J. L., C. A. Reynolds, and S. L. Grove. 1999. Distribution of California Gnatcatchers on Camp Pendleton Marine Corps Base. Prepared for U.S. Marine Corps, Oceanside, California (Contract No. M00681-97-C-0035). Unpublished technical report, February 14, Manomet Center for Conservation Sciences, MA.
- Atwood, J.L. and J.S. Bolsinger. 1992. Elevational distribution of California Gnatcatchers in the United States. *Journal of Field Ornithology* 63: 159-168.
- Atwood, J.L. and D.R. Bontrager. 2001. California Gnatcatcher (*Polioptila californica*). In A. Poole and F. Gill (eds.) *The Birds of North America* No. 574. Philadelphia, PA.
- Bontrager, D. 1991. Habitat Requirements, Home Range and Breeding Biology of the California Gnatcatcher (*Polioptila californica*) in South Orange County. Santa Margarita Company.
- United States Fish and Wildlife Service. 1991. Summary of the proposed rule to list the coastal California gnatcatcher (*Polioptila californica*) as endangered in California and Baja, Mexico. September. 114 pp.
- United States Fish and Wildlife Service. 1993. Federal Register, Endangered and Threatened Wildlife and Plants; Determination of Threatened Status for the Coastal California Gnatcatcher; Final Rule. March 30.
- United States Fish and Wildlife Service. 1997 Coastal California Gnatcatcher (*Polioptila californica californica*) Presence/Absence Survey Protocol. Available at: [https://www.fws.gov/ventura/docs/species/protocols/cagn/coastal-gnatcatcher\\_survey-guidelines.pdf](https://www.fws.gov/ventura/docs/species/protocols/cagn/coastal-gnatcatcher_survey-guidelines.pdf).
- United States Fish and Wildlife Service. 2007. Federal Register, Endangered and Threatened Wildlife and Plants; Revised Designation of Critical Habitat for the Coastal California Gnatcatcher (*Polioptila californica californica*); Final Rule. December 19.
- Unitt, P. A. 2004. *San Diego County Bird Atlas*. Proceedings of the San Diego Society of Natural History, No. 39. San Diego Natural History Museum

# Attachment 1

---

15-Day Notification of Intent Letter





**Rincon Consultants, Inc.**

2215 Faraday Avenue, Suite A  
Carlsbad, California 92008

760 918 9444

info@rinconconsultants.com  
www.rinconconsultants.com

April 18, 2023

Ms. Stacey Love  
United States Fish and Wildlife Service  
Carlsbad Fish and Wildlife Office  
2177 Salk Avenue, Suite 250  
Carlsbad, California 92008  
Via email: [stacey\\_love@fws.gov](mailto:stacey_love@fws.gov)

**Subject: Notification of Intent to Conduct Protocol Breeding Season Surveys for Coastal California Gnatcatcher for the Woodward Specific Plan Development Project in San Marcos, San Diego County, California.**

Dear Ms. Love:

In accordance with the current United States (U.S.) Fish and Wildlife Service (USFWS) survey protocol for coastal California gnatcatcher (*Polioptila californica californica*; CAGN) (dated February 28, 1997), please consider this letter as notification of the intent to commence protocol breeding season surveys for the Woodward Specific Plan Development Project (project site). The 8.57-acre project site is located to the east of Woodward Street and north of East Mission Road and is situated 0.4 mile north of Highway 78. The project site is located within the U.S. Geological Survey 7.5-minute topographic quadrangle *San Marcos, California* (See attached map, Figure 1).

The project is located within the boundaries of the San Diego County Multiple Habitat Conservation Program (MHCP) subarea plan and is located adjacent to a 7.73-acre open space MHCP Hardline Reserve land to the north, east, and west of the project site across Woodward Street. The surveys would be conducted for 15.48 acres of Diegan coastal sage scrub and disturbed Diegan coastal sage scrub habitat within the project site and adjacent to the north, east south and west.

The survey area is outside of the County of San Diego's Natural Community Conservation Plan, therefore a minimum of six (6) surveys will be conducted at least one (1) week apart from May 3 through June 30 following the 15-day notification period. A final report will be submitted to the USFWS within 45 days following the field surveys adhering to the report guidelines outlined in the USFWS Coastal California Gnatcatcher Presence/Absence Survey Protocol.

All CAGN surveys will be conducted by Kelly Rios, 10(a)(1)(A) permit no. TE 018909-5 with Rincon Consultants, Inc. (Rincon) within the 15.48-acre survey area.

Please let us know if you have any questions or concerns.

Sincerely,

**Rincon Consultants, Inc.**

A handwritten signature in black ink, appearing to read "Jared Reed", is written over a light blue horizontal line.


Jared Reed, Senior Biologist/Project Manager

## **Attachments**

Figure 1 USGS Topographic Map

Figure 2 Project Location

Figure 1 Project Site



Basemap provided by National Geographic Society, Esri, and their licensors © 2023. San Marcos Quadrangle. T12S R03W S11.  
The topographic representation depicted in this map may not portray all of the features currently found in the vicinity today and/or features depicted in this map may have changed since the original topographic map was assembled.



Figure 2 Project Location

