
		Appendix
Riological Re	sources Sunna	orting Information
Biological ite.	sources suppr	



ity of Anaheim—The Mill Residential Townhome Project (DEV2023-000 nitial Study/Mitigated Negative Declaration	
	D.1. Chariel status Charies Table
	B.1 - Special-status Species Table




Table 1: Special-status Plant Species Evaluated

Scientific Name	Status				
Common Name	USFWS ¹	CDFW ²	CNPS ³	Habitat Description ⁴	Potential to Occur and Rationale ⁵
Abronia villosa var. aurita chaparral sand-verbena	_	_	1B.1	Annual herb found in chaparral, coastal scrub, desert dune habitat. Grows in sandy soils. Elevation: 75 – 1600 m Blooming period: (January) March – September	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
<i>Aphanisma blitoides</i> Aphanisma	_	_	1B.2	Annual herb found in coastal bluff scrub, coastal dunes, coastal scrub habitat. Sometimes grows in gravelly or sandy soils. Elevation: 1 – 305 m. Blooming Period: February – June	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
<i>Astragalus hornii</i> var. <i>hornii</i> Horn's milk-vetch	_	_	1B.1	Annual herb found in alkali sink, meadows, seeps, playas, wetland-riparian, and lake margins, often in alkaline soils. Elevation: 60 – 850 meters Blooming period: May – October	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Astragalus pycnostachyus var. Ianosissimus Ventura Marsh milk-vetch	FE	SE	1B.1	Perennial herb found in marshes and swamps, coastal dunes, coastal scrub. Within reach of high tide or protected by barrier beaches, more rarely near seeps on sandy bluffs. Elevation: 1 – 35 m. Blooming period: (June)August – October	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Atriplex coulteri Coulter's saltbush	_	_	1B.2	Perennial herb found in valley and foothill grassland, coastal bluff scrub, coastal dunes, coastal scrub habitat. Sometimes grows in clay alkaline soils. Elevation: 3 – 460 m. Blooming Period: March – October	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Atriplex pacifica south coast saltscale	_	_	1B.1	Annual herb found in chenopod scrub, playas, and vernal pool habitat. Grows in alkaline soils. Elevation: 25-1900 m. Blooming period: June – October	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Atriplex parishii Parish's brittlescale	_	_	1B.1	Annual herb found in vernal pools, chenopod scrub, playas. Usually on drying alkali flats with fine soils. Elevation: 25 – 1900 m. Blooming period: June – October	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.

Scientific Name		Status			
Common Name	USFWS ¹	CDFW ²	CNPS ³	Habitat Description⁴	Potential to Occur and Rationale ⁵
Atriplex serenana var. davidsonii Davidson's saltscale	_	_	1B.2	Annual herb found in coastal bluff scrub and coastal scrub habitat. Grows in alkaline soils. Elevation: 10-200 m. Blooming period: April – October	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Calochortus weedii var. intermedius intermediate mariposa-lily	_	_	1B.2	Perennial bulbiferous herb found valley and foothill grassland, chaparral, coastal scrub habitat. Grows in rocky soils. Elevation: May – July Blooming period: 105 – 855 m.	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Calystegia felix lucky morning-glory	_	_	1B.1	Annual rhizomatous herb found in alkaline meadows and seeps, riparian alluvial scrub habitat. Elevation: 30 – 215 m. Blooming period: March – September	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Camissoniopsis lewisii Lewis' evening-primrose	_	_	3	Annual herb found in valley and foothill grassland, coastal bluff scrub, cismontane woodland, coastal dunes, coastal scrub habitat. Sometimes grows in sandy or clay soils. Elevation: 0 – 300 m. Blooming period: March – May(June)	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Centromadia parryi ssp. australis southern tarplant	_	_	1B.1	Annual herb found in margins of marshes and swamps, valley and foothill grassland, vernal pool habitat. Grows in vernally mesic soils. Elevation: 0 – 480 m. Blooming period: May – November	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Chloropyron maritimum ssp. maritimum salt marsh bird's-beak	FE	SE	1B.2	Hemiparasitic herb found in coastal dunes, coastal salt marshes and swamp habitat. Elevation: 0 – 30 m. Blooming period: May – October(November)	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Dudleya multicaulis many-stemmed dudleya	_	_	1B.2	Perennial herb found in chaparral, valley and foothill grassland, coastal scrub habitat. Often grows in clay soils. Elevation: 15 – 790 m. Blooming period: April – July	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Eriastrum densifolium ssp. sanctorum Santa Ana River woollystar	FE	SE	18.1	Perennial herb found in chaparral, coastal alluvial fan scrub habitat. Sometimes grows in gravelly or sandy soils. Elevation: 91 – 610 m. Blooming period: April - September	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.

Scientific Name		Status			
Common Name	USFWS ¹	CDFW ²	CNPS ³	Habitat Description ⁴	Potential to Occur and Rationale ⁵
Eryngium aristulatum var. parishii San Diego button-celery	FE	SE	1B.1	Annual/perennial herb found in coastal scrub, valley and foothill grassland, vernal pool habitat. Grows in mesic soils. Elevation: 20 – 620 m. Blooming period: April – June	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Hordeum intercedens vernal barley	_	_	3.2	Annual herb found in coastal dunes, coastal scrub, valley and foothill grassland, vernal pool habitat. Grows in depressions or saline flats. Elevation: 5 – 1000 m. Blooming period: March – June	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Isocoma menziesii var. decumbens decumbent goldenbush	_	_	1B.2	Perennial shrub found in chaparral, coastal scrub habitat. Often grows in disturbed, sandy soils. Elevation: 10 – 250 m. Blooming period: April – November	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Lasthenia glabrata ssp. coulteri Coulter's goldfields	_	_	1B.1	Annual herb found in coastal salt marshes, playas, vernal pool habitat. Usually found on alkaline soils in playas, sinks, and grasslands. Elevation: 1 – 1375 m. Blooming period: February – June	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Nama stenocarpa mud nama	_	_	2B.2	Annual /perennial herb found in marsh and swamp habitat. Grows on lake margins and riverbanks. Elevation: 5 – 500 m. Blooming period: January – July	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Nasturtium gambelii Gambel's water cress	FE	ST	18.1	Perennial rhizomatous herb found in brackish freshwater marsh and swamp habitat. Elevation: 5 – 330 m. Blooming period: April – October	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Navarretia prostrata prostrate vernal pool navarretia	_	_	1B.2	Annual herb found in coastal scrub, meadows and seeps, valley and foothill grassland, vernal pool habitat. Grows in alkaline and mesic soils. Elevation: 3 – 1210 m. Blooming period: April – July	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.

Colombisio Name		Status			
Scientific Name Common Name	USFWS ¹	CDFW ²	CNPS ³	Habitat Description ⁴	Potential to Occur and Rationale ⁵
Nemacaulis denudata var. denudate coast woolly-heads	_	_	1B.2	Annual herb found in coastal dunes. Elevation: 0 – 100 m. Blooming period: April – September	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Orcuttia californica California Orcutt grass	FE	CE	1B.1	Annual herb found in vernal pool habitat. Elevation: 15 – 660 m. Blooming period: April – August	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Pentachaeta aurea ssp. allenii Allen's pentachaeta	_	_	18.1	Annual herb found in coastal scrub openings, valley and foothill grassland habitat. Elevation: 75 – 520 m. Blooming period: March – June	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Phacelia ramosissima var. austrolitoralis Phacelia ramosissima var. austrolitoralis	_	_	3.2	Perennial herb found in chaparral, coastal dunes, coastal scrub, coastal salt marsh and swamp habitat. Sometimes grows in rocky and sandy soils. Elevation: 5 – 300 m. Blooming period: March – August	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Phacelia stellaris Brand's star phacelia	_	_	1B.1	Annual herb found in coastal scrub, coastal dunes. Open areas. Elevation: 3 – 400 m. Blooming period: March – June	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Sagittaria sanfordii Sanford's arrowhead	_	_	1B.2	Emergent perennial rhizomatous herb found in shallow freshwater marsh and swamp habitat. Elevation: 0 – 650 m. Blooming period: May – October(November)	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Senecio aphanactis chaparral ragwort	_	_	2B.2	Annual herb found in chaparral, cismontane woodland, coastal scrub habitat. Sometimes grows in alkaline soils. Elevation: 15 – 800 m. Blooming period: January – April(May)	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Sidalcea neomexicana salt spring checkerbloom	_	_	2B.2	Found in playas, chaparral, coastal scrub, lower montane coniferous forest, Mojave Desert scrub, alkali springs and marshes. Elevation: 0 – 1,530 m. Bloom period: March – June	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.

= Delisted in accordance with the FESA.

= Federally Proposed to be Delisted.

= Not federally listed

MBTA = protected by the Migratory Bird Treaty Act

Rank 3 = Plants about which we need more information—

Rank 4 = Plants of limited distribution—A Watch List

Blooming period: Months in parentheses are uncommon.

A Review List

Scientific Name		Status			
Common Name	USFWS ¹	CDFW ²	CNPS ³	Habitat Description⁴	Potential to Occur and Rationale ⁵
Suaeda esteroa estuary seablite	_	_	1B.2	Perennial herb found in coastal salt marshes and swamps. Elevation: 0 – 5 m. Blooming period: (January – May)July – October	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
Symphyotrichum defoliatum San Bernardino aster	_	_	1B.2	Found in meadows and seeps, marshes and swamps, coastal scrub, cismontane woodland, lower montane coniferous forest, grassland. Prefers vernally mesic grassland or near ditches, streams, and springs; especially disturbed areas. Elevation: 2 – 2,040 m. Bloom period: July – November	None. The project site is entirely developed and does not contain suitable natural vegetation communities and substates for this plant to grow. Therefore, this species is not expected to occur on-site.
				Code Designations	
¹ Federal Status:	¹ Federal Status: 2024 USFWS Listing			² State Status: 2024 CDFW Listing	³ CNPS: 2024 CNPS Listing
ESU = Evolutionary Significant Unit is a distinctive population.				SE = Listed as endangered under the CESA. ST = Listed as threatened under the CESA.	Rank 1A = Plants species that presumed extinct in California.
FE = Listed as endangered under the FESA. FT = Listed as threatened under the FESA.				SSC = Species of Special Concern as identified by the CDFW. FP = Listed as fully protected under FGC.	Rank 1B = Plant species that are rare, threatened, or endangered in California and elsewhere.
FC = Candidate for listing under FESA.	(threatene	d or endan	gered)	CFG = FGC =protected by FGC 3503.5 CR = Rare in California.	Rank 2 = Plant species that are rare, threatened, or endangered in California, but more common
FD = Delisted in accordan	ce with the	FFSΔ			elsewhere.

Not state listed

Sources:

California Department of Fish and Wildlife (CDFW). 2024. CNDDB RareFind 5 California Natural Diversity Database Query for Special-Status Species. Website: https://map.dfg.ca.gov/rarefind/view/RareFind.aspx. Accessed January 11, 2024.

California Native Plant Society (CNPS). 2024. California Native Plant Society Rare and Endangered Plant Inventory. Website: http://www.rareplants.cnps.org/. Accessed January 11, 2024. California Department of Fish and Wildlife (CDFW). 2024. Biogeographic Information and Observation System (BIOS 6). Website: https://map.dfg.ca.gov/bios/. Accessed January 11, 2024.

⁴ Habitat Description: Habitat description adapted from CNDDB and CNPS online inventory or other specified source.

⁵ Potential to Occur and Rationale: Location of recorded species occurrences determined by geospatial information from BIOS 6 or other specified source*.

Table 2: Special-status Wildlife Species Evaluated

Scientific Name		tus								
Common Name	USFWS ¹	CDFW ²	Habitat Description ³	Potential to Occur and Rationale ⁴						
Birds	3irds									
Accipiter cooperii Cooper's hawk	— MBTA	FGC WL	Occurs and nests in deciduous and mixed forests and open woodland habitats. Year-round resident in southern California, and tolerant of urban areas with an abundance of trees.	None. There are no recent or historical records within five miles of the project site. The project is entirely developed and does not contain suitable vegetation communities to support this species. Therefore, this species is not expected to occur on-site.						
Agelaius tricolor tricolored blackbird	— MBTA	FT FGC SSC	Occurs and nests in large freshwater marshes with dense stands of hydrophytic vegetation (cattails, bulrushes, etc.). Short-distance migrant.	None. There are no recent or historical records within five miles of the project site. The project site does not contain marsh habitat to support this species. Therefore, this species is not expected to occur on-site.						
Aimophila ruficeps canescens southern California rufous- crowned sparrow	— MBTA	— WL	Occurs and nests on steep, often rocky hillsides with grass and forb patches in coastal sage and Riversidean alluvial fan sage scrub and sparse mixed chaparral habitats. Year-round resident in southern California.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable grass or scrub habitat to support this species. Therefore, this species is not expected to occur on-site.						
Ammodramus savannarum grasshopper sparrow	— MBTA	_ SSC	Occurs and nests in dense grasslands on rolling hills, lowland plains, in valleys and on hillsides on lower mountain slopes.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable nesting habitat to support this species. Therefore, this species is not expected to occur on-site.						
Asio otus long-eared owl	— MBTA	FP FGC SSC	Occurs and nests in conifer, oak, riparian, pinyon-juniper, and desert woodlands that are either open or are adjacent to grasslands, meadows, or shrublands	None. There is one historical occurrence recorded in 1925, within five miles of the project site. The project site does not contain suitable habitat or mammal burrows to support this species. Therefore, this species is not expected to occur on-site.						
Athene cunicularia burrowing owl	_	FGC SSC	Occurs and nests in open, dry annual or perennial grasslands, deserts, and scrublands characterized by low-growing vegetation. A subterranean nester, dependent upon burrowing mammals, most notably the California ground squirrel. Short-distance migrant.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable habitat or mammal burrows to support this species. Therefore, this species is not expected to occur on-site.						
Buteo regalis ferruginous hawk	— MBTA	— WL	Grassland and arid shrublands with an abundance of prey species, such as pocket gophers, black-tailed jackrabbits, and desert cottontails. Will winter near cultivated fields that have an abundance of pocket gophers.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable foraging habitat to support this species. Therefore, this species is not expected to occur on-site.						

Scientific Name	Stat	tus		
Common Name	USFWS ¹	CDFW ²	Habitat Description ³	Potential to Occur and Rationale ⁴
Buteo swainsoni Swainson's hawk (nesting)	— MBTA	ST CFG	Occurs and nests in grasslands with scattered trees, juniper-sage flats, riparian areas, savannahs, and agricultural or ranch lands with groves or lines of trees. Requires adjacent suitable foraging areas such as grasslands, or alfalfa or grain fields supporting rodent populations. Long-distance migrant.	None. There is one historical occurrence recorded in 1888, within five miles of the project site. The project site lacks suitable foraging and nesting habitat. This species is believed to be locally extirpated and therefore, this species is not expected to occur on-site.
Campylorhynchus brunneicapillus sandiegensis coastal cactus wren	MBTA CONCCP HCP	_ SSC	Occurs and nests in coastal sage scrub. Requires tall opuntia cactus or <i>Yucca</i> trees for nesting and roosting.	None. There is one historical occurrence recorded in 2002, within five miles of the project site. The project site does not contain suitable sage scrub vegetation to support this species. Therefore, this species is not expected to occur on-site.
Charadrius nivosus nivosus western snowy plover	FT MBTA	_ SSC	Occurs in sandy beaches, salt pond levees and shores of large alkali lakes. Needs sandy, gravelly or friable soils for nesting.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable marsh or aquatic habitat to support this species. Therefore, this species is not expected to occur on-site.
Coccyzus americanus occidentalis western yellow-billed cuckoo	FT MBTA	SE	Found in riparian forest along the broad lower flood-bottoms of larger river systems. Nests in riparian jungles of willow, often mixed with cottonwoods; understory consists of blackberry, nettles, and wild grape.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable riparian habitat for this species to nest. Therefore, this species is not expected to occur on-site.
Coturnicops noveboracensis yellow rail	— MBTA	FGC SSC	Occurs in shallow marshes and wet meadows. In winter, this species inhabits drier fresh-water and brackish marshes, as well as dense, deep grass, and rice fields.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable aquatic habitat or marsh vegetation to support this species. Therefore, this species is not expected to occur on-site.
Elanus leucurus white-tailed kite	MBTA FP	FGC	Occurs and nests in grasslands and open coastal scrub in coastal and valley lowlands; rarely found away from agricultural areas. Inhabits herbaceous, open stages of most habitats mostly in cismontane California. Year-round resident in southern California.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable riparian scrub vegetation to support this species. Therefore, this species is not expected to occur on-site.
Eremophila alpestris actia California horned lark	— MBTA	FGC WL	Occurs and nests in open areas with sparse vegetation. Year-round resident in southern California.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable bare or sparsely vegetated areas free from human disturbance for this species to nest. Therefore, this species is not expected to occur on-site.

Scientific Name	Stat	us		
Common Name	USFWS ¹	CDFW ²	Habitat Description ³	Potential to Occur and Rationale ⁴
Falco peregrinus anatum American peregrine falcon	FD MBTA CONCCP HCP	SD FP	Occurs near bodies of water, including wetlands, lakes, and rivers, in open areas with cliffs, ledges, and canyons nearby for cover and nesting. They will also nest on banks, dunes, and mounds. Their nests are a scrape on a depression or ledge on an open site. They will nest on human-made structures, and occasionally uses tree or snag cavities or old nests of other raptors.	None. There are no recent or historical records within five miles of the project site. The project site does not contain cliffs or tall man-made structures suitable for nesting. Therefore, this species is not expected to occur on-site.
Icteria virens yellow-breasted chat	— MBTA	FGC SSC	Occurs and nests in riparian thickets of willow and other bushy tangles near watercourses. Long-distance migrant.	None. There is one historical occurrence recorded in 2001, within five miles of the project site. The project site does not contain suitable riparian habitat for this species to nest. Therefore, this species is not expected to occur on-site.
Laterallus jamaicensis coturniculus California black rail	— MBTA	ST FP	Occurs and nests in freshwater marshes, wet meadows, and shallow margins of saltwater marshes bordering larger bays. Needs water depths of about 1 inch that do not fluctuate during the year and dense vegetation for nesting habitat.	None. There are no recent or historical records within five miles of the project site. The project site lacks marsh or meadow habitat to support this species. Therefore, this species is not expected to occur on-site.
Passerculus sandwichensis beldingi Belding's savannah sparrow	— MBTA	SE	Occurs in coastal wetlands and marshes.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable wetland or marsh habitat for this species. Therefore, this species is not expected to occur on-site.
Polioptila californica californica coastal California gnatcatcher	FT MBTA	_ SSC	Obligate, permanent resident of coastal sage scrub below 2,500 ft in Southern California. Often found in arid washes, on mesas and slopes. Not all areas classified as coastal sage scrub are occupied.	Low. There are three historical occurrences recorded within five miles of the project site, the most recent in 2007. The project site does not contain suitable coastal sage scrub habitat to support this species. Therefore, this species is not expected to occur onsite.
Rallus obsoletus levipes light-footed Ridgway's rail	FE FP	SE	Occurs in salt marshes traversed by tidal sloughs, where cordgrass and pickleweed (required for nesting or escape cover) are the dominant vegetation.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable salt marsh habitat for this species to nest. Therefore, this species is not expected to occur on-site.
Riparia riparia bank swallow	— MBTA	ST	Nests in riparian scrub and riparian woodland. Requires vertical banks/cliffs with fine-textured/sandy soils near streams, rivers, lakes, ocean to dig nesting hole.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable riparian habitat for this species to nest. Therefore, this species is not expected to occur on-site.

Scientific Name	Scientific Name Status			
Common Name	USFWS ¹	CDFW ²	Habitat Description ³	Potential to Occur and Rationale ⁴
Rynchops niger black skimmer	— MBTA	_ SSC	Occurs and nests on gravel bars, low islets, and sandy beaches, in unvegetated sites.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable bare or sparsely vegetated areas free from human disturbance for this species to nest. Therefore, this species is not expected to occur on-site.
Setophaga petechia yellow warbler	— MBTA	FGC SSC	Occurs and nests in willow shrubs and thickets, cottonwoods, sycamores, ash, and alders, predominantly in riparian habitats. Long-distance migrant.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable riparian habitat for this species to nest. Therefore, this species is not expected to occur on-site.
Sternula antillarum browni California least tern	FE MBTA	SE FP	Occurs and nests on bare or sparsely vegetated, flat substrates: sand beaches, alkali flats, landfills, or paved areas.	Low. There are two historical occurrences recorded within five miles of the project site, the most recent in 2018. The project site does not contain suitable bare or sparsely vegetated areas free from human disturbance for this species to nest. Therefore, this species is not expected to occur on-site.
Vireo bellii pusillus least Bell's vireo	FE MBTA CONCCP HCP	SE CONCCP HCP	Summer resident of southern California in low riparian in vicinity of water or in dry river bottoms; below 2000 ft. Nests placed along margins of bushes or on twigs projecting into pathways, usually willow, <i>Baccharis</i> , mesquite.	None. There is one historical occurrence recorded within five miles of the project site, the most recent in 2013. The project site does not contain suitable riparian habitat for this species to nest. Therefore, this species is not expected to occur on-site.
Fish				
Catostomus santaanae Santa Ana sucker	FT	_	Occurs in south coastal streams in the Los Angeles basin. Prefers sand-rubble-boulder bottoms, cool, clear water, and algae.	None. There are no recent or historical records within five miles of the project site. The project site is fully developed and lacks suitable aquatic habitat to support this species. Therefore, this species is not expected to occur on-site.
Oncorhynchus mykiss irideus (pop. 10) steelhead (southern California DPS¹)	FE	SE	Occurs from the Santa Maria River to the Tijuana River at the United States and Mexican Border in seasonally accessible rivers and streams. Steelhead require cool, clean water and appropriate aggregate for spawning. Southern steelhead likely have greater physiological tolerances to warmer water and more variable conditions.	None. There are no recent or historical records within five miles of the project site. The project site is fully developed and lacks suitable aquatic habitat to support this species. Therefore, this species is not expected to occur on-site.

¹ Federal listing refers to populations from Santa Maria River south to southern extent of range (San Mateo Creek in San Diego County).

Scientific Name Common Name	Stat	tus		Potential to Occur and Rationale ⁴
	USFWS ¹	CDFW ²	Habitat Description ³	
Amphibians				
Spea hammondii western spadefoot	FT CONCCP HCP	_ SSC	Occurs in open areas with sandy or gravelly soils in mixed woodlands, grasslands, coastal sage and Riversidean alluvial fan sage scrub, chaparral, sandy washes, lowlands, river floodplains, alluvial fans, playas, alkali flats, foothills, and mountains. Breeds in ephemeral rain pools that do not contain bullfrogs, fish, or crayfish.	None. There are no recent or historical records within five miles of the project site. The project site is fully developed and lacks suitable aquatic habitat to support this species. Therefore, this species is not expected to occur on-site.
Invertebrates				
Bombus crotchii Crotch bumble bee	_	SC	This species occurs primarily in California, including coastal habitats, western Mojave Desert, San Joaquin Valley, and adjacent foothills through most of southwestern California. It inhabits arid grasslands and shrublands, and its food sources including milkweeds, pincushions, lupines, clovers, phacelias, sages, clarkias, poppies, and buckwheat.	Low. There are no recent or historical records within five miles of the project site. The project site does contain a few marginally suitable ornamental plants which this species may forage on. However, more suitable foraging habitat can be found in undeveloped areas nearby.
Danaus plexippus plexippus pop. 1 monarch - California overwintering population	FC	_	Occurs in temperate climates, such as eastern and western North America and undergoes long-distance migration. Lays eggs on obligate milkweed host plant (primarily <i>Asclepias spp.</i>)	None. There are no recent or historical records within five miles of the project site. The project site does not contain obligatory milkweed host plants to support this species. Therefore, this species is not expected to occur on-site.
Euphydryas editha quino quino checkerspot butterfly	FE	_	Found in sunny openings within chaparral & coastal sage shrublands in parts of Riverside & San Diego counties. Hills and mesas near the coast. This species needs high densities of food plants <i>Plantago erecta, P. insularis,</i> and <i>Orthocarpus purpurescens</i> .	None. There is one historical occurrence recorded within five miles of the project site, the most recent in 1930. The project site does not contain suitable chaparral or sage scrub or food plants to support this species. Therefore, this species is not expected to occur on-site.
Crustaceans				
Branchinecta sandiegonensis San Diego fairy shrimp	FE	_	Small, shallow (5-30 cm deep) vernal pools with cool water (10-20°C), moderate alkalinity and conductivity, and less than 1 m deep.	None. There are no recent or historical records within five miles of the project site. The project site is fully developed and lacks suitable vernal pool habitat to support this species. Therefore, this species is not expected to occur on-site.
Streptocephalus woottoni Riverside fairy shrimp	FE	_	Vernal pools on the Santa Rosa Plateau on Murrieta stony clay loams, Las Posas series, Wyman clay loam, and Willows soils.	None. There are no recent or historical records within five miles of the project site. The project site is fully developed and lacks suitable vernal pool habitat to support this species. Therefore, this species is not expected to occur on-site.

Scientific Name	Status						
Common Name	USFWS ¹	CDFW ²	Habitat Description ³	Potential to Occur and Rationale ⁴			
Mammals							
Choeronycteris mexicana Mexican long-tongued bat	_	– SSC	Occurs in pinon and juniper woodlands, riparian scrub, and Sonoran thorn woodland. Feeds on nectar and pollen of night-blooming succulents. Roosts in relatively well-lit caves, and in and around buildings.	None. There are no recent or historical records within five miles of the project site. No suitable woodland or roosting habitat in the form of high vacant buildings are present on-site. Therefore, this species is not expected to occur on-site.			
Eumops perotis californicus western mastiff bat	_	– SSC	Occurs in many open, semi-arid to arid habitats, including conifer and deciduous woodlands, coastal scrub, grasslands, chaparral. Roosts in crevices in cliff faces, high buildings, trees, and tunnels.	Low. There are three historical occurrences recorded within five miles of the project site, the most recent in 1990. No suitable roosting habitat in the form of high vacant buildings are present on-site. Therefore, this species is not expected to occur on-site.			
Lasiurus xanthinus western yellow bat	_	_ SSC	Occurs in valley foothill riparian, desert riparian, desert wash, and palm oasis habitats. Roosts in skirts of dead fronds in both native and non-native palm trees.	None. There are no recent or historical records within five miles of the project site. Marginally suitable roosting habitat in the form of ornamental trees are present on-site. However, the project site lacks nearby water sources and is entirely developed subject to human disturbance.			
Microtus californicus stephensi south coast marsh vole	_	– SSC	Occurs in tidal marshes in Los Angeles, Orange and southern Ventura counties.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable salt marsh habitat for this species. Therefore, this species is not expected to occur on-site.			
Nyctinomops femorosaccus pocketed free-tailed bat	_	_ SSC	Occurs in pine-juniper woodlands, desert scrub, palm oasis, desert wash, and desert riparian habitats. Roosts in caves, crevices, mines, tunnels, and man-made structures.	None. There are no recent or historical records within five miles of the project site. Marginally suitable roosting habitat in the form of ornamental trees and man-made structures are present on-site. However, the project site lacks nearby water sources and is entirely developed subject to human disturbance.			
Nyctinomops macrotis big free-tailed bat	_	_ SSC	Occurs in rocky habitats in arid landscapes and has been found in a variety of plant associations, including desert shrub, woodlands, and evergreen forests. Roosts by day in crevices on cliff faces. These bats forage mostly for large moths, but may eat other insects as well.	Low. There are no recent or historical records within five miles of the project site. Suitable roosting habitat in the form of ornamental trees and man-made structures are present on-site. However, the project site lacks nearby water sources and is entirely developed subject to human disturbance.			
Perognathus longimembris pacificus Pacific pocket mouse	FE CONCCP HCP	– SSC CONCCP HCP	Occurs in coastal strand, coastal dunes, and coastal sage scrub growing on marine terraces. Prefers soils of fine alluvial sands near the ocean.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable coastal habitat for this species. Therefore, this species is not expected to occur on-site.			

Scientific Name	Status			
Common Name	USFWS ¹	CDFW ²	Habitat Description ³	Potential to Occur and Rationale ⁴
Sorex ornatus salicornicus southern California saltmarsh shrew	-	_ ssc	Occurs in coastal marshes in Los Angeles, Orange and Ventura counties. Requires dense vegetation and woody debris for cover.	None. There are no recent or historical records within five miles of the project site. The project site does not contain suitable coastal marsh habitat for this species. Therefore, this species is not expected to occur on-site.
<i>Taxidea taxus</i> American badger	-	_ ssc	Occurs in drier open stages of most shrub, forest, and herbaceous habitats with friable soils. Requires sufficient food sources (rodents), friable soils, and open, uncultivated ground. Digs large burrows.	None. There are no recent or historical records within five miles of the project site. The project is entirely developed and does not contain suitable vegetation communities to support this species. Therefore, this species is not expected to occur on-site.
Reptiles				
Anniella stebbinsi Southern California legless lizard	_	– SSC	Occurs south of the Transverse Range, extending to northwestern Baja California with disjunct populations in the Tehachapi and Piute Mountains in Kern County. Occurs in a variety of habitats but often sandy or loose loamy soils under sparse vegetation. They prefer soils with a high moisture content.	Low. There are two historical occurrences recorded within five miles of the project site, the most recent in 2019. The project is entirely developed and does not contain suitable vegetation communities to support this species. Therefore, this species is not expected to occur on-site.
Aspidoscelis hyperythra orange-throated whiptail	— CONCCP HCP	— WL	Occurs primarily on coarse soils in open coastal sage and Riversidean alluvial fan sage scrub habitat.	None. There are no recent or historical records within five miles of the project site. The project is entirely developed and does not contain suitable vegetation communities to support this species. Therefore, this species is not expected to occur on-site.
Aspidoscelis tigris stejnegeri coastal whiptail	-	_ SSC	Occurs in dry, open areas with sparse foliage in coastal sage and Riversidean alluvial fan sage scrub, chaparral, woodland, and riparian habitats.	None. There are no recent or historical records within five miles of the project site. The project is entirely developed and does not contain suitable vegetation communities to support this species. Therefore, this species is not expected to occur on-site.
Crotalus ruber red-diamond rattlesnake	— CONCCP HCP	_ SSC	Occurs in arid, rocky areas in creosote scrub, coastal sage and Riversidean alluvial fan sage scrub, chaparral, oak and pine woodlands, grasslands, on cultivated areas.	None. There are no recent or historical records within five miles of the project site. The project is entirely developed and does not contain suitable vegetation communities to support this species. Therefore, this species is not expected to occur on-site.
Emys marmorata western pond turtle	-	_ SSC	Occurs in ponds, lakes, rivers, streams, creeks, marshes, and irrigation ditches with abundant vegetation, and either rocky or muddy bottoms, in woodland, forest, and grassland. In streams, prefers pools to shallower areas. Logs, rocks, cattail mats, and exposed banks are required for basking. May enter brackish water and even seawater.	None. There is one recent occurrence within five miles of the project site, the most recent recorded in 1989. The project parcel does not contain suitable aquatic to support this species. Therefore, this species is not expected to occur on-site.

Scientific Name	Status					
		CDFW ²	Habitat Description ³	Potential to Occur and Rationale ⁴		
Phrynosoma blainvillii coast horned lizard	_	SSC	Occurs in open areas with sandy soil and low vegetation in grasslands, coniferous forests, woodlands, and chaparral.	None. There are two historical occurrences recorded within five miles of the project site, the most recent in 1971. The project is entirely developed and does not contain suitable vegetation communities to support this species. Therefore, this species is not expected to occur on-site.		
Salvadora hexalepis virgultea coast patch-nosed snake	_	_ SSC	Occurs in brushy or shrubby vegetation. Dependent on small mammal burrows.	None. There are no recent or historical records within five miles of the project site. The project is entirely developed and does not contain suitable vegetation communities to support this species. Therefore, this species is not expected to occur on-site.		

Code Designations

	¹ Federal Status: 2024 USFWS Listing	² State Status: 2024 CDFW Listing
ESU	= Evolutionary Significant Unit is a distinctive population.	SE = Listed as endangered under the CESA.
FE	= Listed as endangered under the FESA.	ST = Listed as threatened under the CESA.
FT	= Listed as threatened under the FESA.	SSC = Species of Special Concern as identified by the CDFW.
FC	= Candidate for listing (threatened or endangered) under FESA.	FP = Listed as fully protected under FGC.
FD	= Delisted in accordance with the FESA.	CFG = FGC =protected by FGC 3503.5
FPD	= Federally Proposed to be Delisted.	SC = Candidate endangered under the CESA.
MBTA	= protected by the Migratory Bird Treaty Act	WL = Species monitored by CDFW "Watch List."
-	= Not federally listed	— = Not state listed


Habitat Description: Habitat description adapted from CNDDB or other specified source*.

Sources:


California Department of Fish and Wildlife (CDFW). 2024. CNDDB RareFind 5 California Natural Diversity Database Query for Special-Status Species. Website: https://map.dfg.ca.gov/rarefind/view/RareFind.aspx. Accessed January 11, 2024.

California Department of Fish and Wildlife (CDFW). 2024. Biogeographic Information and Observation System (BIOS 6). Website: https://map.dfg.ca.gov/bios/. Accessed January 11, 2024.

⁴ Potential to Occur and Rationale: Location of recorded species occurrences determined by geospatial information from BIOS 6 or other specified source*.



Initial Study/Mitigated Negative Declaration	mome Project (DEV2023-00042) on
	D. 2. California Natural Discovita Database Communica
	B.2 - California Natural Diversity Database Summary





California Department of Fish and Wildlife California Natural Diversity Database



Query Criteria:

Quad IS (Anaheim (3311778) OR Los Alamitos (3311871) OR Crange (3311777) OR Whittier (3311881) OR La Habra (3311788) OR Yorba Linda (3311787) OR Seal Beach (3311861) OR Newport Beach (3311768) OR Tustin (3311767))

Species	Element Code	Federal Status	State Status	Global Rank	State Rank	Rare Plant Rank/CDFW SSC or FP
Abronia villosa var. aurita	PDNYC010P1	None None	None Status	G5T2?	State Rank	1B.1
chaparral sand-verbena	1 2111 00 101 1	TTOTIC	None	00121	OL.	10.1
Accipiter cooperii	ABNKC12040	None	None	G5	S4	WL
Cooper's hawk						
Agelaius tricolor	ABPBXB0020	None	Threatened	G1G2	S2	SSC
tricolored blackbird						
Aimophila ruficeps canescens	ABPBX91091	None	None	G5T3	S4	WL
southern California rufous-crowned sparrow						
Ammodramus savannarum	ABPBXA0020	None	None	G5	S3	SSC
grasshopper sparrow						
Anniella stebbinsi	ARACC01060	None	None	G3	S3	SSC
Southern California legless lizard						
Aphanisma blitoides	PDCHE02010	None	None	G3G4	S2	1B.2
aphanisma						
Ardea herodias	ABNGA04010	None	None	G5	S4	
great blue heron						
Asio otus	ABNSB13010	None	None	G5	S3?	SSC
long-eared owl						
Aspidoscelis hyperythra	ARACJ02060	None	None	G5	S2S3	WL
orange-throated whiptail						
Aspidoscelis tigris stejnegeri	ARACJ02143	None	None	G5T5	S3	SSC
coastal whiptail						
Astragalus hornii var. hornii	PDFAB0F421	None	None	GUT1	S1	1B.1
Horn's milk-vetch						
Astragalus pycnostachyus var. Ianosissimus Ventura Marsh milk-vetch	PDFAB0F7B1	Endangered	Endangered	G2T1	S1	1B.1
Athene cunicularia	ABNSB10010	None	None	G4	S2	SSC
burrowing owl						
Atriplex coulteri	PDCHE040E0	None	None	G3	S1S2	1B.2
Coulter's saltbush						
Atriplex pacifica south coast saltscale	PDCHE041C0	None	None	G4	S2	1B.2
Atriplex parishii	PDCHE041D0	None	None	G1G2	S1	1B.1
Parish's brittlescale						
Atriplex serenana var. davidsonii	PDCHE041T1	None	None	G5T1	S1	1B.2
Davidson's saltscale						
Bombus crotchii	IIHYM24480	None	Candidate	G2	S2	
Crotch bumble bee			Endangered			





Smaaira	Fla	Fada::-101 1	04-4- 04-4	Class D	State D	Rare Plant Rank/CDFW
Species	Element Code	Federal Status	State Status	Global Rank	State Rank	SSC or FP
Bombus pensylvanicus American bumble bee	IIHYM24260	None	None	G3G4	S2	
	LODD A 00000	Endonment	Niere	00	04	
Branchinecta sandiegonensis	ICBRA03060	Endangered	None	G2	S1	
San Diego fairy shrimp	A DAUGO (04.00	Nama	Nama	C4	0004	10/1
Buteo regalis ferruginous hawk	ABNKC19120	None	None	G4	S3S4	WL
Buteo swainsoni	A DNIZ C 10070	None	Throatonod	CE	S4	
Swainson's hawk	ABNKC19070	None	Threatened	G5	34	
California Walnut Woodland	CTT71210CA	None	None	G2	S2.1	
California Walnut Woodland California Walnut Woodland	C1171210CA	None	None	G2	32.1	
Calochortus plummerae	PMLIL0D150	None	None	G4	S4	4.2
Plummer's mariposa-lily	I WEILOD 130	None	None	04	04	4.2
Calochortus weedii var. intermedius	PMLIL0D1J1	None	None	G3G4T3	S3	1B.2
intermediate mariposa-lily	1 WEIEOD 101	None	None	000410	00	10.2
Calystegia felix	PDCON040P0	None	None	G1Q	S1	1B.1
lucky morning-glory	1 2001101010	TTOTIC	140110	O T Q	01	15.1
Campylorhynchus brunneicapillus sandiegensis	ABPBG02095	None	None	G5T3Q	S2	SSC
coastal cactus wren	7.5. 5002000	110110	110110	30104	02	
Catostomus santaanae	AFCJC02190	Threatened	None	G1	S1	
Santa Ana sucker						
Centromadia parryi ssp. australis	PDAST4R0P4	None	None	G3T2	S2	1B.1
southern tarplant						
Charadrius nivosus nivosus	ABNNB03031	Threatened	None	G3T3	S3	SSC
western snowy plover						
Chelonia mydas	ARAAA02010	Threatened	None	G3	S1	
green turtle						
Chloropyron maritimum ssp. maritimum	PDSCR0J0C2	Endangered	Endangered	G4?T1	S1	1B.2
salt marsh bird's-beak						
Choeronycteris mexicana	AMACB02010	None	None	G3G4	S1	SSC
Mexican long-tongued bat						
Cicindela hirticollis gravida	IICOL02101	None	None	G5T2	S2	
sandy beach tiger beetle						
Cicindela latesignata	IICOL02110	None	None	G2G3	S1	
western beach tiger beetle						
Cicindela senilis frosti	IICOL02121	None	None	G2G3T1T3	S1	
senile tiger beetle						
Coccyzus americanus occidentalis	ABNRB02022	Threatened	Endangered	G5T2T3	S1	
western yellow-billed cuckoo						
Coelus globosus	IICOL4A010	None	None	G1G2	S1S2	
globose dune beetle						
Coturnicops noveboracensis	ABNME01010	None	None	G4	S2	SSC
yellow rail						





Species	Element Code	Federal Status	State Status	Global Rank	State Rank	Rare Plant Rank/CDFW SSC or FP
Species Crotalus ruber	ARADE02090		None Status	G4 G4	State Rank	SSC OF FP
red-diamond rattlesnake	ARADE02090	None	None	G4	33	330
Danaus plexippus plexippus pop. 1	IILEPP2012	Candidate	None	G4T1T2Q	S2	
monarch - California overwintering population	IILLFF2012	Candidate	None	G41112Q	32	
Dudleya multicaulis	PDCRA040H0	None	None	G2	S2	1B.2
many-stemmed dudleya	1 0010040110	None	None	02	02	10.2
Elanus leucurus	ABNKC06010	None	None	G5	S3S4	FP
white-tailed kite	7.5141.000010	140110	140110	00	0001	• •
Emys marmorata	ARAAD02030	Proposed	None	G3G4	S3	SSC
western pond turtle	, , , , , , , , , , , , , , , , , , , ,	Threatened	110110	333.		
Eremophila alpestris actia	ABPAT02011	None	None	G5T4Q	S4	WL
California horned lark						
Eriastrum densifolium ssp. sanctorum	PDPLM03035	Endangered	Endangered	G4T1	S1	1B.1
Santa Ana River woollystar		3.	3			
Eryngium aristulatum var. parishii	PDAPI0Z042	Endangered	Endangered	G5T1	S1	1B.1
San Diego button-celery		J	J			
Eumops perotis californicus	AMACD02011	None	None	G4G5T4	S3S4	SSC
western mastiff bat						
Euphydryas editha quino	IILEPK405L	Endangered	None	G4G5T1T2	S1S2	
quino checkerspot butterfly						
Falco peregrinus anatum	ABNKD06071	Delisted	Delisted	G4T4	S3S4	
American peregrine falcon						
Glyptostoma gabrielense	IMGASB1010	None	None	G2	S3	
San Gabriel chestnut						
Habroscelimorpha gabbii	IICOL02080	None	None	G2G4	S1	
western tidal-flat tiger beetle						
Helianthus nuttallii ssp. parishii	PDAST4N102	None	None	G5TX	SX	1A
Los Angeles sunflower						
lcteria virens	ABPBX24010	None	None	G5	S4	SSC
yellow-breasted chat						
lsocoma menziesii var. decumbens	PDAST57091	None	None	G3G5T2T3	S2	1B.2
decumbent goldenbush						
Lasionycteris noctivagans	AMACC02010	None	None	G3G4	S3S4	
silver-haired bat						
Lasiurus cinereus	AMACC05032	None	None	G3G4	S4	
hoary bat						
Lasiurus xanthinus	AMACC05070	None	None	G4G5	S3	SSC
western yellow bat						
Lasthenia glabrata ssp. coulteri	PDAST5L0A1	None	None	G4T2	S2	1B.1
Coulter's goldfields						
Laterallus jamaicensis coturniculus	ABNME03041	None	Threatened	G3T1	S2	FP
California black rail						





s State Status None None None None Threatened None None None Candidate	Global Rank G5T3 G5T2T3 G5 G4G5 G1 G2 G3G4T2 G5	\$tate Rank \$3 \$2 \$4 \$1\$\$2 \$1 \$2 \$2 \$3 \$3 \$3	SSC or FP 4.3 SSC 2B.2 1B.1 1B.2 1B.2 SSC
None None Threatened None None None	G5T2T3 G5 G4G5 G1 G2 G3G4T2 G5	\$2 \$4 \$1\$2 \$1 \$2 \$2 \$3	2B.2 1B.1 1B.2 1B.2
None None Threatened None None None	G5 G4G5 G1 G2 G3G4T2 G5	\$4 \$1\$2 \$1 \$2 \$2 \$3	2B.2 1B.1 1B.2 1B.2
None None Threatened None None None	G5 G4G5 G1 G2 G3G4T2 G5	\$4 \$1\$2 \$1 \$2 \$2 \$3	2B.2 1B.1 1B.2 1B.2
None Threatened None None None	G4G5 G1 G2 G3G4T2 G5	\$1\$2 \$1 \$2 \$2 \$3	1B.1 1B.2 1B.2
None Threatened None None None	G4G5 G1 G2 G3G4T2 G5	\$1\$2 \$1 \$2 \$2 \$3	1B.1 1B.2 1B.2
Threatened None None None None	G1 G2 G3G4T2 G5	\$1 \$2 \$2 \$3	1B.1 1B.2 1B.2
Threatened None None None None	G1 G2 G3G4T2 G5	\$1 \$2 \$2 \$3	1B.1 1B.2 1B.2
None None None	G2 G3G4T2 G5	S2 S2 S3	1B.2 1B.2
None None None	G2 G3G4T2 G5	S2 S2 S3	1B.2 1B.2
None None None	G3G4T2 G5	S2 S3	1B.2
None None None	G3G4T2 G5	S2 S3	1B.2
None None	G5	S3	
None None	G5	S3	
None			SSC
	G5	S3	
	G5	S3	
Candidate			SSC
Candidate			
Endangered	G5T1Q	S1	
Endangered	G1	S1	1B.1
Ū			
None	G5	S4	WL
None	G4	S2	
Endangered	G5T3	S3	
S			
None	G4T1	S1	1B.1
None	G5T2	S2	SSC
	00.2	02	
None	G1	S1	1B.1
Home	0.		15.1
None	G4	S 4	SSC
None	04	04	330
None	G4G5T3O	52	SSC
NONE	0 1 0013Q	52	300
	C2T1T2	C 1	FP
Endangered	G31112	31	T.F.
Endangered		62	
	CF	ა ა	
	None None None Endangered	None G1 None G4 None G4G5T3Q	None G1 S1 None G4 S4 None G4G5T3Q S2 Endangered G3T1T2 S1





Species	Element Code	Federal Status	State Status	Global Rank	State Rank	Rare Plant Rank/CDFW SSC or FP
Rynchops niger	ABNNM14010	None	None	G5	S2	SSC
black skimmer						
Sagittaria sanfordii	PMALI040Q0	None	None	G3	S3	1B.2
Sanford's arrowhead						
Salvadora hexalepis virgultea	ARADB30033	None	None	G5T4	S3	SSC
coast patch-nosed snake						
Senecio aphanactis	PDAST8H060	None	None	G3	S2	2B.2
chaparral ragwort						
Setophaga petechia	ABPBX03010	None	None	G5	S3	SSC
yellow warbler						
Sidalcea neomexicana	PDMAL110J0	None	None	G4	S2	2B.2
salt spring checkerbloom						
Sorex ornatus salicornicus	AMABA01104	None	None	G5T1?	S1	SSC
southern California saltmarsh shrew						
Southern California Arroyo Chub/Santa Ana Sucker Stream	CARE2330CA	None	None	GNR	SNR	
Southern California Arroyo Chub/Santa Ana Sucker Stream						
Southern Coast Live Oak Riparian Forest	CTT61310CA	None	None	G4	S4	
Southern Coast Live Oak Riparian Forest						
Southern Coastal Salt Marsh	CTT52120CA	None	None	G2	S2.1	
Southern Coastal Salt Marsh						
Southern Cottonwood Willow Riparian Forest	CTT61330CA	None	None	G3	S3.2	
Southern Cottonwood Willow Riparian Forest						
Southern Dune Scrub	CTT21330CA	None	None	G1	S1.1	
Southern Dune Scrub						
Southern Foredunes	CTT21230CA	None	None	G2	S2.1	
Southern Foredunes						
Southern Sycamore Alder Riparian Woodland Southern Sycamore Alder Riparian Woodland	CTT62400CA	None	None	G4	S4	
Southern Willow Scrub	CTT63320CA	None	None	G3	S2.1	
Southern Willow Scrub						
Spea hammondii	AAABF02020	Proposed	None	G2G3	S3S4	SSC
western spadefoot		Threatened				
Sternula antillarum browni	ABNNM08103	Endangered	Endangered	G4T2T3Q	S2	FP
California least tern						
Streptocephalus woottoni	ICBRA07010	Endangered	None	G1G2	S2	
Riverside fairy shrimp						
Suaeda esteroa	PDCHE0P0D0	None	None	G3	S2	1B.2
estuary seablite						
Symphyotrichum defoliatum	PDASTE80C0	None	None	G2	S2	1B.2
San Bernardino aster						
Taxidea taxus	AMAJF04010	None	None	G5	S3	SSC
American badger						




California Department of Fish and Wildlife California Natural Diversity Database



Species	Element Code	Federal Status	State Status	Global Rank	State Rank	Rare Plant Rank/CDFW SSC or FP
Trigonoscuta dorothea dorothea	IICOL51021	None	None	G1T1	S1	
Dorothy's El Segundo Dune weevil						
Tryonia imitator mimic tryonia (=California brackishwater snail)	IMGASJ7040	None	None	G2	S2	
Vireo bellii pusillus least Bell's vireo	ABPBW01114	Endangered	Endangered	G5T2	S3	

Record Count: 106

Initial Study/Mitigated Negati	ve Declaration				
B.3 - California N	Native Plant So	ociety Rare I	Plant Invento	ory Database	Summary
Dio Gamerina i		colory mane .		or y Database	y diriinidi y





CNPS Rare Plant Inventory

Search Results

49 matches found. Click on scientific name for details

									CA RARE			
▲ SCIENTIFIC NAME	COMMON NAME	FAMILY	LIFEFORM	BLOOMING PERIOD	FED LIST	STATE LIST	GLOBAL RANK		PLANT RANK		DATE ADDED	РНОТО
<u>Abronia</u> <u>maritima</u>	red sand- verbena	Nyctaginaceae	perennial herb	Feb-Nov	None	None	G4	S3?	4.2		1994- 01-01	©2003
												Christopher L. Christie
	chaparral sand-verbena	Nyctaginaceae	annual herb	(Jan)Mar- Sep	None	None	G5T2?	S2	1B.1		2001- 01-01	© 2011
												Aaron E.
<u>Aphanisma</u> <u>blitoides</u>	aphanisma	Chenopodiaceae	annual herb	Feb-Jun	None	None	G3G4	S 2	1B.2		1980- 01-01	Sims
												© 2010 Larry Sward
<u>Astragalus</u> <u>hornii var. hornii</u>	Horn's milk- vetch	Fabaceae	annual herb	May-Oct	None	None	GUT1	S1	1B.1		2006- 12-01	No Photo Available
	Ventura Marsh milk- vetch	Fabaceae	perennial herb	(Jun)Aug- Oct	FE	CE	G2T1	S1	1B.1	Yes	1974- 01-01	No Photo Available
	Coulter's saltbush	Chenopodiaceae	perennial herb	Mar-Oct	None	None	G3	S1S2	1B.2		1994- 01-01	No Photo Available
	south coast saltscale	Chenopodiaceae	annual herb	Mar-Oct	None	None	G4	S2	1B.2		1994- 01-01	No Photo Available
<u>Atriplex parishii</u>	Parish's brittlescale	Chenopodiaceae	annual herb	Jun-Oct	None	None	G1G2	S1	1B.1		1988- 01-01	No Photo Available
<u>Atriplex</u> serenana var. davidsonii	Davidson's saltscale	Chenopodiaceae	annual herb	Apr-Oct	None	None	G5T1	S1	1B.2		1994- 01-01	No Photo Available

<u>Calochortus</u> <u>catalinae</u>	Catalina mariposa lily	Liliaceae	perennial bulbiferous herb	(Feb)Mar- Jun	None	None	G3G4	S3S4	4.2	Yes	1974- 01-01	No Photo Available
<u>Calochortus</u> <u>plummerae</u>	Plummer's mariposa-lily	Liliaceae	perennial bulbiferous herb	May-Jul	None	None	G4	S4	4.2	Yes	1994- 01-01	No Photo Available
<u>Calochortus</u> <u>weedii var.</u> <u>intermedius</u>	intermediate mariposa-lily	Liliaceae	perennial bulbiferous herb	May-Jul	None	None	G3G4T3	S3	1B.2	Yes	1994- 01-01	No Photo Available
<u>Calystegia felix</u>	lucky morning- glory	Convolvulaceae	annual rhizomatous herb	Mar-Sep	None	None	G1Q	S1	1B.1	Yes	2014- 07-16	No Photo Available
<u>Camissoniopsis</u> <u>lewisii</u>	Lewis' evening- primrose	Onagraceae	annual herb	Mar- May(Jun)	None	None	G4	S4	3		1994- 01-01	No Photo Available
<u>Centromadia</u> <u>parryi ssp.</u> <u>australis</u>	southern tarplant	Asteraceae	annual herb	May-Nov	None	None	G3T2	S2	1B.1		1994- 01-01	No Photo Available
<u>Chloropyron</u> <u>maritimum ssp.</u> <u>maritimum</u>	salt marsh bird's-beak	Orobanchaceae	annual herb (hemiparasitic)	May- Oct(Nov)	FE	CE	G4?T1	S1	1B.2		1974- 01-01	No Photo Available
<u>Cistanthe</u> <u>maritima</u>	seaside cistanthe	Montiaceae	annual herb	(Feb)Mar- Jun(Aug)	None	None	G3G4	S 3	4.2		1980- 01-01	No Photo Available
<u>Convolvulus</u> <u>simulans</u>	small- flowered morning- glory	Convolvulaceae	annual herb	Mar-Jul	None	None	G4	S4	4.2		1994- 01-01	No Photo Available
<u>Deinandra</u> paniculata	paniculate tarplant	Asteraceae	annual herb	(Mar)Apr- Nov	None	None	G4	S4	4.2		2001- 01-01	No Photo Available
<u>Dudleya</u> multicaulis	many- stemmed dudleya	Crassulaceae	perennial herb	Apr-Jul	None	None	G2	S2	1B.2	Yes	1974- 01-01	No Photo Available
<u>Eleocharis</u> parvula	small spikerush	Cyperaceae	perennial herb	(Apr)Jun- Aug(Sep)	None	None	G5	S3	4.3		1980- 01-01	©2018 Ron Vanderhoff
Eriastrum densifolium ssp. sanctorum	Santa Ana River woollystar	Polemoniaceae	perennial herb	Apr-Sep	FE	CE	G4T1	S1	1B.1	Yes	1980- 01-01	No Photo Available
<u>Eryngium</u> aristulatum var. parishii	San Diego button-celery	Apiaceae	annual/perennial herb	Apr-Jun	FE	CE	G5T1	S1	1B.1		1974- 01-01	No Photo Available
Helianthus nuttallii ssp. parishii	Los Angeles sunflower	Asteraceae	perennial rhizomatous herb	Aug-Oct	None	None	G5TX	SX	1A	Yes	1974- 01-01	No Photo Available

<u>Hordeum</u> <u>intercedens</u>	vernal barley	Poaceae	annual herb	Mar-Jun	None	None	G3G4	S3S4	3.2		1994- 01-01	No Photo Available
Isocoma menziesii var. decumbens	decumbent goldenbush	Asteraceae	perennial shrub	Apr-Nov	None	None	G3G5T2T3	S2	1B.2		1994- 01-01	No Photo Available
Juglans californica	Southern California black walnut	Juglandaceae	perennial deciduous tree	Mar-Aug	None	None	G4	S4	4.2	Yes	1994- 01-01	© 2020 Zoya
Juncus acutus ssp. leopoldii	southwestern spiny rush	Juncaceae	perennial rhizomatous herb	(Mar)May- Jun	None	None	G5T5	S4	4.2		1988- 01-01	Akulova © 2019 Belinda Lo
<u>Lasthenia</u> g <u>labrata ssp.</u> coulteri	Coulter's goldfields	Asteraceae	annual herb	Feb-Jun	None	None	G4T2	S2	1B.1		1994- 01-01	© 2013
<u>Lepidium</u> virginicum var. robinsonii	Robinson's pepper-grass	Brassicaceae	annual herb	Jan-Jul	None	None	G5T3	S3	4.3		1994- 01-01	© 2015 Keir Morse
<u>Lycium</u> <u>californicum</u>	California box-thorn	Solanaceae	perennial shrub	Mar- Aug(Dec)	None	None	G4	S4	4.2		2001- 01-01	No Photo Available
Microseris douglasii ssp. platycarpha	small- flowered microseris	Asteraceae	annual herb	Mar-May	None	None	G4T4	S4	4.2		2001- 01-01	© 2015 Richard Spellenberg
<u>Nama</u> <u>stenocarpa</u>	mud nama	Namaceae	annual/perennial herb	Jan-Jul	None	None	G4G5	S1S2	2B.2		1994- 01-01	No Photo Available
<u>Nasturtium</u> g <u>ambelii</u>	Gambel's water cress	Brassicaceae	perennial rhizomatous herb	Apr-Oct	FE	СТ	G1	S1	1B.1		1980- 01-01	No Photo Available
<u>Navarretia</u> <u>prostrata</u>	prostrate vernal pool navarretia	Polemoniaceae	annual herb	Apr-Jul	None	None	G2	S2	1B.2	Yes	2001- 01-01	No Photo Available
Nemacaulis denudata var. denudata	coast woolly- heads	Polygonaceae	annual herb	Apr-Sep	None	None	G3G4T2	S2	1B.2		1994- 01-01	No Photo Available
<u>Orcuttia</u> <u>californica</u>	California Orcutt grass	Poaceae	annual herb	Apr-Aug	FE	CE	G1	S1	1B.1		1974- 01-01	No Photo Available


<u>Pentachaeta</u> <u>aurea ssp.</u> allenii	Allen's pentachaeta	Asteraceae	annual herb	Mar-Jun	None	None	G4T1	S1	1B.1	Yes	2008- 05-08	©2008 Bob
<u>Phacelia hubbyi</u>	Hubby's	Hydrophyllaceae	annual herh	Apr-Jul	None	None	G4	S4	4.2	Yes	2007-	Allen
rnacetta nace	phacelia	. iyaropiiyiiaccac	armaar norb	7 lp. 7 di	110110		0.			.05	02-02	No Photo
												Available
<u>Phacelia</u> 	south coast	Hydrophyllaceae	perennial herb	Mar-Aug	None	None	G5?T3Q	S3	3.2		2007-	
<u>ramosissima var.</u> <u>austrolitoralis</u>	phacelia										05-17	No Photo Available
Phacelia stellaris	•	Hydrophyllaceae	annual herh	Mar-Jun	None	None	G1	S1	1B.1		1994-	Available
rnacella stellaris	phacelia	Пуспорпупасеае	annual nerb	IVIAI -JUII	None	None	G1	31	10.1		01-01	No Photo
												Available
<u>Quercus</u>	Engelmann	Fagaceae	perennial	Mar-Jun	None	None	G3	S3	4.2		1988-	
<u>engelmannii</u>	oak		deciduous tree								01-01	No Photo
												Available
<u>Romneya</u> <u>coulteri</u>	Coulter's matilija	Papaveraceae	perennial rhizomatous	Mar- Jul(Aug)	None	None	G4	S4	4.2		1974- 01-01	No Photo
counter	рорру		herb	Jui(Aug)							01 01	Available
<u>Sagittaria</u> <u>sanfordii</u>	Sanford's arrowhead	Alismataceae	perennial rhizomatous herb (emergent)	May- Oct(Nov)	None	None	G3	S3	1B.2	Yes	1984- 01-01	
												©2013
												Debra L. Cook
Canacia	chanarral	Actoração	annual herb	lan	None	None	C2	S2	2B.2		1994-	COOK
<u>Senecio</u> <u>aphanactis</u>	chaparral ragwort	Asteraceae	annuai nerb	Jan- Apr(May)	None	None	d5	32	ZD.Z		01-01	No Photo
_,	3			1								Available
<u>Sidalcea</u>	salt spring	Malvaceae	perennial herb	Mar-Jun	None	None	G4	S2	2B.2		1994-	
<u>neomexicana</u>	checkerbloom										01-01	No Photo
												Available
<u>Suaeda esteroa</u>	estuary seablite	Chenopodiaceae	perennial herb	(Jan- May)Jul-	None	None	G3	S2	1B.2		1984- 01-01	No Photo
	seabilte			Oct							01-01	Available
Suaeda taxifolia	woolly	Chenopodiaceae	perennial	Jan-Dec	None	None	G4	S4	4.2		1994-	
	seablite	,	evergreen shrub								01-01	No Photo
												Available
<u>Symphyotrichum</u>		Asteraceae	perennial	Jul-Nov	None	None	G2	S2	1B.2	Yes	2004-	
<u>defoliatum</u>	Bernardino aster		rhizomatous herb								01-01	No Photo Available

Showing 1 to 49 of 49 entries

Suggested Citation:

California Native Plant Society, Rare Plant Program. 2024. Rare Plant Inventory (online edition, v9.5). Website https://www.rareplants.cnps.org [accessed 11 January 2024].

City of Anaheim—The Mill Residential Townhome Project (DEV2023-00042) Initial Study/Mitigated Negative Declaration	
	B.4 - IpaC Report
	Dr. Ipac Reperc



Thi

hal

(US

bel

tha

X

al

'S

t

iced

[Pac recource list

On Thursday, January 11th(Today), the ability to generate Official Species Lists and Determination Key Documents will be unavailable for a short time beginning at 17:00 EST. There may also be a disruption to some of the map layers. These functions will be restored as soon as they are available. We appreciate your patience with this temporary disruption.

However, determining the likelihood and extent of effects a project may have on trust resources typically requires gathering additional site-specific (e.g., vegetation/species surveys) and project-specific (e.g., magnitude and timing of proposed activities) information.

Below is a summary of the project information you provided and contact information for the USFWS office(s) with jurisdiction in the defined project area. Please read the introduction to each section that follows (Endangered Species, Migratory Birds, USFWS Facilities, and NWI Wetlands) for additional information applicable to the trust resources addressed in that section.

Location

Orange County, California



Local office

Carlsbad Fish And Wildlife Office

\((760) 431-9440

(760) 431-5901



Endangered species

This resource list is for informational purposes only and does not constitute an analysis of project level impacts.

The primary information used to generate this list is the known or expected range of each species. Additional areas of influence (AOI) for species are also considered. An AOI includes areas outside of the species range if the species could be indirectly affected by activities in that area (e.g., placing a dam upstream of a fish population even if that fish does not occur at the dam site, may indirectly impact the species by reducing or eliminating water flow downstream). Because species can move, and site conditions can change, the species on this list are not guaranteed to be found on or near the project area. To fully determine any potential effects to species, additional site-specific and project-specific information is often required.

Section 7 of the Endangered Species Act **requires** Federal agencies to "request of the Secretary information whether any species which is listed or proposed to be listed may be present in the area of such proposed action" for any project that is conducted, permitted, funded, or licensed by any Federal agency. A letter from the local office and a species list which fulfills this requirement can **only** be obtained by requesting an official species list from either the Regulatory Review section in IPaC (see directions below) or from the local field office directly.

For project evaluations that require USFWS concurrence/review, please return to the IPaC website and request an official species list by doing the following:

- 1. Draw the project location and click CONTINUE.
- 2. Click DEFINE PROJECT.
- 3. Log in (if directed to do so).
- 4. Provide a name and description for your project.
- 5. Click REQUEST SPECIES LIST.

Listed species¹ and their critical habitats are managed by the <u>Ecological Services Program</u> of the U.S. Fish and Wildlife Service (USFWS) and the fisheries division of the National Oceanic and Atmospheric Administration (NOAA Fisheries²).

Species and critical habitats under the sole responsibility of NOAA Fisheries are **not** shown on this list. Please contact <u>NOAA Fisheries</u> for <u>species under their jurisdiction</u>.

1. Species listed under the <u>Endangered Species Act</u> are threatened or endangered; IPaC also shows species that are candidates, or proposed, for listing. See the <u>listing status page</u> for more information. IPaC only shows species that are regulated by USFWS (see FAQ).

2. <u>NOAA Fisheries</u>, also known as the National Marine Fisheries Service (NMFS), is an office of the National Oceanic and Atmospheric Administration within the Department of Commerce.

Endangered

Endangered

The following species are potentially affected by activities in this location:

Mammals

NAME STATUS

Pacific Pocket Mouse Perognathus longimembris pacificus

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/8080

Stephens' Kangaroo Rat Dipodomys stephensi (incl. D. **Threatened**

cascus)

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/3495

Birds

NAME STATUS

California Least Tern Sternula antillarum browni

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/8104

California Spotted Owl Strix occidentalis occidentalis Proposed Endangered

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/7266

Coastal California Gnatcatcher Polioptila californica Threatened

californica

Wherever found

There is **final** critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/8178

Hawaiian Petrel Pterodroma sandwichensis Endangered

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/6746

Least Bell's Vireo Vireo bellii pusillus

Wherever found

There is **final** critical habitat for this species. Your location does not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/5945

Light-footed Ridgeway""s Rail Rallus obsoletus levipes

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/6035

Short-tailed Albatross Phoebastria (=Diomedea) albatrus

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/433

Southwestern Willow Flycatcher Empidonax traillii extimus

Wherever found

There is **final** critical habitat for this species. Your location does not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/6749

Western Snowy Plover Charadrius nivosus nivosus

There is **final** critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/8035

Endangered

Endangered

Endangered

Endangered

Threatened

Reptiles

NAME STATUS

Southwestern Pond Turtle Actinemys pallida

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/4768

Proposed Threatened

Amphibians

NAME STATUS

Arroyo (=arroyo Southwestern) Toad Anaxyrus californicus

Wherever found

There is **final** critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/3762

Endangered

Fishes

NAME STATUS

Santa Ana Sucker Catostomus santaanae

There is **final** critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/3785

Threatened

Tidewater Goby Eucyclogobius newberryi

Wherever found

There is **final** critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/57

Endangered

Insects

NAME STATUS

Monarch Butterfly Danaus plexippus

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/9743

Candidate

Quino Checkerspot Butterfly Euphydryas editha quino (=E.

e. wrighti)

Wherever found

There is final critical habitat for this species. Your location does not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/5900

Endangered

Crustaceans

NAME STATUS

Riverside Fairy Shrimp Streptocephalus woottoni

Wherever found

There is **final** critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/8148

San Diego Fairy Shrimp Branchinecta sandiegonensis

Wherever found

There is **final** critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/6945

Endangered

Endangered

Vernal Pool Fairy Shrimp Branchinecta lynchi

Wherever found

There is final critical habitat for this species. Your location does not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/498

Threatened

Flowering Plants

NAME **STATUS**

Big-leaved Crownbeard Verbesina dissita

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/8049

Threatened

Braunton's Milk-vetch Astragalus brauntonii

Wherever found

There is final critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/5674

Endangered

Encinitas Baccharis Baccharis vanessae

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/3343

Threatened

Laguna Beach Liveforever Dudleya stolonifera

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/7919

Threatened

Munz's Onion Allium munzii

Wherever found

There is **final** critical habitat for this species. Your location does not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/2951

Salt Marsh Bird's-beak Cordylanthus maritimus ssp.

maritimus

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/6447

San Diego Ambrosia Ambrosia pumila

Wherever found

There is final critical habitat for this species. Your location does not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/8287

San Diego Button-celery Eryngium aristulatum var. parishii

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/5937

San Diego Thornmint Acanthomintha ilieffolia

Wherever found

There is **final** critical habitat for this species. Your location does not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/351

Santa Monica Mountains Dudleyea Dudleya cymosa ssp.

ovatifolia

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/2538

Slender-horned Spineflower Dodecahema leptoceras

Wherever found

No critical habitat has been designated for this species.

https://ecos.fws.gov/ecp/species/4007

Endangered

Endangered

Endangered

Endangered

Threatened

Threatened

Endangered

Spreading Navarretia Navarretia fossalis

Wherever found

There is **final** critical habitat for this species. Your location does not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/1334

Thread-leaved Brodiaea Brodiaea filifolia

Wherever found

There is **final** critical habitat for this species. Your location overlaps the critical habitat.

https://ecos.fws.gov/ecp/species/6087

Ventura Marsh Milk-vetch Astragalus pycnostachyus var.

lanosissimus Wherever found

> There is final critical habitat for this species. Your location does SULT not overlap the critical habitat.

https://ecos.fws.gov/ecp/species/1160

Threatened

Threatened

Endangered

Final

Critical habitats

Potential effects to critical habitat(s) in this location must be analyzed along with the endangered species themselves.

This location overlaps the critical habitat for the following species:

NAME **TYPF**

Arroyo (=arroyo Southwestern) Toad Anaxyrus californicus Final

https://ecos.fws.gov/ecp/species/3762#crithab

Braunton's Milk-vetch Astragalus brauntonii

https://ecos.fws.gov/ecp/species/5674#crithab

Coastal California Gnatcatcher Polioptila californica Final

californica

https://ecos.fws.gov/ecp/species/8178#crithab

Riverside Fairy Shrimp Streptocephalus woottoni Final

https://ecos.fws.gov/ecp/species/8148#crithab

Final San Diego Fairy Shrimp Branchinecta sandiegonensis

https://ecos.fws.gov/ecp/species/6945#crithab

Santa Ana Sucker Catostomus santaanae
https://ecos.fws.gov/ecp/species/3785#crithab

Thread-leaved Brodiaea Brodiaea filifolia
https://ecos.fws.gov/ecp/species/6087#crithab

Tidewater Goby Eucyclogobius newberryi
https://ecos.fws.gov/ecp/species/57#crithab

Western Snowy Plover Charadrius nivosus nivosus
https://ecos.fws.gov/ecp/species/8035#crithab

Bald & Golden Eagles

There are no documented cases of eagles being present at this location. However, if you believe eagles may be using your site, please reach out to the local Fish and Wildlife Service office.

Additional information can be found using the following links:

- Eagle Management https://www.fws.gov/program/eagle-management
- Measures for avoiding and minimizing impacts to birds
 https://www.fws.gov/library/collections/avoiding-and-minimizing-incidental-take-migratory-birds
- Nationwide conservation measures for birds
 https://www.fws.gov/sites/default/files/documents/nationwide-standard-conservation-measures.pdf
- Supplemental Information for Migratory Birds and Eagles in IPaC https://www.fws.gov/media/supplemental-information-migratory-birds-and-bald-and-golden-eagles-may-occur-project-action

Bald and Golden Eagle information is not available at this time

What does IPaC use to generate the potential presence of bald and golden eagles in my specified location?

The potential for eagle presence is derived from data provided by the <u>Avian Knowledge Network (AKN)</u>. The AKN data is based on a growing collection of <u>survey</u>, <u>banding</u>, <u>and citizen science datasets</u> and is queried and filtered to return a list of those birds reported as occurring in the 10km grid cell(s) which your project

intersects, and that have been identified as warranting special attention because they are a BCC species in that area, an eagle (<u>Eagle Act</u> requirements may apply). To see a list of all birds potentially present in your project area, please visit the <u>Rapid Avian Information Locator (RAIL) Tool</u>.

What does IPaC use to generate the probability of presence graphs of bald and golden eagles in my specified location?

The Migratory Bird Resource List is comprised of USFWS <u>Birds of Conservation Concern (BCC)</u> and other species that may warrant special attention in your project location.

The migratory bird list generated for your project is derived from data provided by the <u>Avian Knowledge Network (AKN)</u>. The AKN data is based on a growing collection of <u>survey</u>, <u>banding</u>, <u>and citizen science datasets</u> and is queried and filtered to return a list of those birds reported as occurring in the 10km grid cell(s) which your project intersects, and that have been identified as warranting special attention because they are a BCC species in that area, an eagle (<u>Eagle Act</u> requirements may apply), or a species that has a particular vulnerability to offshore activities or development.

Again, the Migratory Bird Resource list includes only a subset of birds that may occur in your project area. It is not representative of all birds that may occur in your project area. To get a list of all birds potentially present in your project area, please visit the <u>Rapid Avian Information Locator (RAIL) Tool</u>.

What if I have eagles on my list?

If your project has the potential to disturb or kill eagles, you may need to obtain a permit to avoid violating the <u>Eagle Act</u> should such impacts occur. Please contact your local Fish and Wildlife Service Field Office if you have questions.

Migratory birds

Certain birds are protected under the Migratory Bird Treaty Act¹ and the Bald and Golden Eagle Protection Act².

Any person or organization who plans or conducts activities that may result in impacts to migratory birds, eagles, and their habitats should follow appropriate regulations and consider implementing appropriate conservation measures, as described below.

- 1. The Migratory Birds Treaty Act of 1918.
- 2. The Bald and Golden Eagle Protection Act of 1940.

Additional information can be found using the following links:

- Eagle Management https://www.fws.gov/program/eagle-management
- Measures for avoiding and minimizing impacts to birds
 <u>https://www.fws.gov/library/collections/avoiding-and-minimizing-incidental-take-migratory-birds</u>
- Nationwide conservation measures for birds https://www.fws.gov/sites/default/files/documents/nationwide-standard-conservation-measures.pdf

 Supplemental Information for Migratory Birds and Eagles in IPaC https://www.fws.gov/media/supplemental-information-migratory-birds-and-bald-and-golden-eagles-may-occur-project-action

Migratory bird information is not available at this time

Tell me more about conservation measures I can implement to avoid or minimize impacts to migratory birds.

Nationwide Conservation Measures describes measures that can help avoid and minimize impacts to all birds at any location year round. Implementation of these measures is particularly important when birds are most likely to occur in the project area. When birds may be breeding in the area, identifying the locations of any active nests and avoiding their destruction is a very helpful impact minimization measure. To see when birds are most likely to occur and be breeding in your project area, view the Probability of Presence Summary. Additional measures or permits may be advisable depending on the type of activity you are conducting and the type of infrastructure or bird species present on your project site.

What does IPaC use to generate the list of migratory birds that potentially occur in my specified location?

The Migratory Bird Resource List is comprised of USFWS <u>Birds of Conservation Concern (BCC)</u> and other species that may warrant special attention in your project location.

The migratory bird list generated for your project is derived from data provided by the <u>Avian Knowledge Network (AKN)</u>. The AKN data is based on a growing collection of <u>survey</u>, <u>banding</u>, <u>and citizen science datasets</u> and is queried and filtered to return a list of those birds reported as occurring in the 10km grid cell(s) which your project intersects, and that have been identified as warranting special attention because they are a BCC species in that area, an eagle (<u>Eagle Act</u> requirements may apply), or a species that has a particular vulnerability to offshore activities or development.

Again, the Migratory Bird Resource list includes only a subset of birds that may occur in your project area. It is not representative of all birds that may occur in your project area. To get a list of all birds potentially present in your project area, please visit the <u>Rapid Avian Information Locator (RAIL) Tool</u>.

What does IPaC use to generate the probability of presence graphs for the migratory birds potentially occurring in my specified location?

The probability of presence graphs associated with your migratory bird list are based on data provided by the <u>Avian Knowledge Network (AKN)</u>. This data is derived from a growing collection of <u>survey</u>, <u>banding</u>, <u>and citizen science datasets</u>.

Probability of presence data is continuously being updated as new and better information becomes available. To learn more about how the probability of presence graphs are produced and how to interpret them, go the Probability of Presence Summary and then click on the "Tell me about these graphs" link.

How do I know if a bird is breeding, wintering or migrating in my area?

To see what part of a particular bird's range your project area falls within (i.e. breeding, wintering, migrating or year-round), you may query your location using the RAIL Tool and look at the range maps provided for birds in your area at the bottom of the profiles provided for each bird in your results. If a bird on your migratory bird species list has a breeding season associated with it, if that bird does occur in your project area, there may be nests present at some point within the timeframe specified. If "Breeds elsewhere" is indicated, then the bird likely does not breed in your project area.

What are the levels of concern for migratory birds?

Migratory birds delivered through IPaC fall into the following distinct categories of concern:

- 1. "BCC Rangewide" birds are <u>Birds of Conservation Concern</u> (BCC) that are of concern throughout their range anywhere within the USA (including Hawaii, the Pacific Islands, Puerto Rico, and the Virgin Islands);
- 2. "BCC BCR" birds are BCCs that are of concern only in particular Bird Conservation Regions (BCRs) in the continental USA; and
- 3. "Non-BCC Vulnerable" birds are not BCC species in your project area, but appear on your list either because of the <u>Eagle Act</u> requirements (for eagles) or (for non-eagles) potential susceptibilities in offshore areas from certain types of development or activities (e.g. offshore energy development or longline fishing).

Although it is important to try to avoid and minimize impacts to all birds, efforts should be made, in particular, to avoid and minimize impacts to the birds on this list, especially eagles and BCC species of rangewide concern. For more information on conservation measures you can implement to help avoid and minimize migratory bird impacts and requirements for eagles, please see the FAQs for these topics.

Details about birds that are potentially affected by offshore projects

For additional details about the relative occurrence and abundance of both individual bird species and groups of bird species within your project area off the Atlantic Coast, please visit the <u>Northeast Ocean Data Portal</u>. The Portal also offers data and information about other taxa besides birds that may be helpful to you in your project review. Alternately, you may download the bird model results files underlying the portal maps through the <u>NOAA NCCOS Integrative Statistical Modeling and Predictive Mapping of Marine Bird Distributions and Abundance on the Atlantic Outer Continental Shelf project webpage.</u>

Bird tracking data can also provide additional details about occurrence and habitat use throughout the year, including migration. Models relying on survey data may not include this information. For additional information on marine bird tracking data, see the <u>Diving Bird Study</u> and the <u>nanotag studies</u> or contact <u>Caleb Spiegel</u> or <u>Pam Loring</u>.

What if I have eagles on my list?

If your project has the potential to disturb or kill eagles, you may need to <u>obtain a permit</u> to avoid violating the Eagle Act should such impacts occur.

Proper Interpretation and Use of Your Migratory Bird Report

The migratory bird list generated is not a list of all birds in your project area, only a subset of birds of priority concern. To learn more about how your list is generated, and see options for identifying what other birds may be in your project area, please see the FAQ "What does IPaC use to generate the migratory birds potentially occurring in my specified location". Please be aware this report provides the "probability of

presence" of birds within the 10 km grid cell(s) that overlap your project; not your exact project footprint. On the graphs provided, please also look carefully at the survey effort (indicated by the black vertical bar) and for the existence of the "no data" indicator (a red horizontal bar). A high survey effort is the key component. If the survey effort is high, then the probability of presence score can be viewed as more dependable. In contrast, a low survey effort bar or no data bar means a lack of data and, therefore, a lack of certainty about presence of the species. This list is not perfect; it is simply a starting point for identifying what birds of concern have the potential to be in your project area, when they might be there, and if they might be breeding (which means nests might be present). The list helps you know what to look for to confirm presence, and helps guide you in knowing when to implement conservation measures to avoid or minimize potential impacts from your project activities, should presence be confirmed. To learn more about conservation measures, visit the FAQ "Tell me about conservation measures I can implement to avoid or minimize impacts to migratory birds" at the bottom of your migratory bird trust resources page.

Facilities

National Wildlife Refuge lands

Any activity proposed on lands managed by the <u>National Wildlife Refuge</u> system must undergo a 'Compatibility Determination' conducted by the Refuge. Please contact the individual Refuges to discuss any questions or concerns.

This location overlaps the following National Wildlife Refuge lands:

LAND ACRES

SEAL BEACH NATIONAL WILDLIFE REFUGE 982.23 acres

Fish hatcheries

There are no fish hatcheries at this location.

Wetlands in the National Wetlands Inventory (NWI)

Impacts to <u>NWI wetlands</u> and other aquatic habitats may be subject to regulation under Section 404 of the Clean Water Act, or other State/Federal statutes.

For more information please contact the Regulatory Program of the local <u>U.S. Army Corps of Engineers District</u>.

Wetland information is not available at this time

This can happen when the National Wetlands Inventory (NWI) map service is unavailable, or for very large projects that intersect many wetland areas. Try again, or visit the <u>NWI map</u> to view wetlands at this location.

Data limitations

The Service's objective of mapping wetlands and deepwater habitats is to produce reconnaissance level information on the location, type and size of these resources. The maps are prepared from the analysis of high altitude imagery. Wetlands are identified based on vegetation, visible hydrology and geography. A margin of error is inherent in the use of imagery; thus, detailed on-the-ground inspection of any particular site may result in revision of the wetland boundaries or classification established through image analysis.

The accuracy of image interpretation depends on the quality of the imagery, the experience of the image analysts, the amount and quality of the collateral data and the amount of ground truth verification work conducted. Metadata should be consulted to determine the date of the source imagery used and any mapping problems.

Wetlands or other mapped features may have changed since the date of the imagery or field work. There may be occasional differences in polygon boundaries or classifications between the information depicted on the map and the actual conditions on site.

Data exclusions

Certain wetland habitats are excluded from the National mapping program because of the limitations of aerial imagery as the pirmary data source used to detect wetlands. These habitats include seagrasses or submerged equatic vegetation that are found in the intertidal and subtidal zones of estuaries and nearshore coastal waters. Some deepwater reef communities (coral or tuberficid worm reefs) have also been excluded from the inventory. These habitats, because of their depth, go undetected by aerial imagery.

Data precautions

Federal, state, and local regulatory agencies with jurisdiction over wetlands may define and describe wetlands in a different manner than that used in this inventory. There is no attempt, in either the design or products of this inventory, to define the limits of proprietary jurisdiction of any Federal, state, or local government or to establish the geographical scope of the regulatory programs of government agencies. Persons intending to engage in activities involving modifications within or adjacent to wetland areas should seek the advice of appropriate Federal, state, or local agencies concerning specified agency regulatory programs and proprietary jurisdictions that may affect such activities.

